

=> fil reg

FILE 'REGISTRY' ENTERED AT 07:39:32 ON 13 DEC 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 12 DEC 2007 HIGHEST RN 957825-32-0

DICTIONARY FILE UPDATES: 12 DEC 2007 HIGHEST RN 957825-32-0

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH June 29, 2007


Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=> d sta que 18

L1 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

VAR G5=H/AK

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 41

CONNECT IS M1 RC AT 45

DEFAULT MLEVEL IS ATOM


DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RSPEC 1
NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L2 2639 SEA FILE=REGISTRY CSS FUL L1
L3 STR




VAR G1=C/38/29
VAR G2=AK/OH/33/36
VAR G3=C/38
VAR G4=C/38/29/42/44
VAR G5=H/AK
NODE ATTRIBUTES:
CONNECT IS M1 RC AT 41
CONNECT IS M1 RC AT 45
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RSPEC 1
NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L5 134 SEA FILE=REGISTRY SUB=L2 CSS FUL L3
L8 131 SEA FILE=REGISTRY ABB=ON PLU=ON L5 NOT (T/ELS OR 14C#)

=> d sta que l18
L1 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

VAR G5=H/AK

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 41

CONNECT IS M1 RC AT 45

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:


RSPEC 1

NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L2 2639 SEA FILE=REGISTRY CSS FUL L1

L9 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

VAR G5=H/AK
 NODE ATTRIBUTES:
 CONNECT IS M1 RC AT 45
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RSPEC 1
 NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE


L11 351 SEA FILE=REGISTRY SUB=L2 CSS FUL L9
 L12 23 SEA FILE=REGISTRY ABB=ON PLU=ON L11 AND NC>=2
 L13 328 SEA FILE=REGISTRY ABB=ON PLU=ON L11 NOT L12
 L14 295 SEA FILE=REGISTRY ABB=ON PLU=ON L13 NOT ((D OR T)/ELS OR
 11C# OR 13C# OR 14C# OR C11# OR C13# OR C14# OR LABELED)
 L15 12 SEA FILE=REGISTRY ABB=ON PLU=ON L14 AND IDS/CI
 L16 283 SEA FILE=REGISTRY ABB=ON PLU=ON L14 NOT L15
 L17 3 SEA FILE=REGISTRY ABB=ON PLU=ON L16 AND NR>=7
 L18 280 SEA FILE=REGISTRY ABB=ON PLU=ON L16 NOT L17

=> d sta que 138

L38 4 SEA FILE=REGISTRY ABB=ON PLU=ON 126-18-1 OR 470-03-1 OR
 16653-88-6 OR 126-19-2

=> d sta que 126

L1 STR




VAR G1=C/38/29
 VAR G2=AK/OH/33/36
 VAR G3=C/38
 VAR G4=C/38/29/42/44
 VAR G5=H/AK
 NODE ATTRIBUTES:
 CONNECT IS M1 RC AT 41
 CONNECT IS M1 RC AT 45
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RSPEC 1
NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L2 2639 SEA FILE=REGISTRY CSS FUL L1
L21 STR



VAR G1=C/38/29
VAR G2=AK/OH/33/36
VAR G3=C/38
VAR G4=C/38/29/42/44
VAR G5=H/AK
VAR G6=46/52/60
VAR G7=AK/CB/50/51
REP G8=(0-1) O
REP G9=(0-1) C
NODE ATTRIBUTES:
CONNECT IS M1 RC AT 45
CONNECT IS M1 RC AT 51
CONNECT IS M1 RC AT 57
CONNECT IS M1 RC AT 58
CONNECT IS M1 RC AT 59
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED
ECOUNT IS M1 O AT 57

GRAPH ATTRIBUTES:


RSPEC 1
NUMBER OF NODES IS 58

STEREO ATTRIBUTES: NONE

L23 324 SEA FILE=REGISTRY SUB=L2 CSS FUL L21
L24 12 SEA FILE=REGISTRY ABB=ON PLU=ON L23 AND NC>=2
L25 64 SEA FILE=REGISTRY ABB=ON PLU=ON L23 AND NR>=7
L26 52 SEA FILE=REGISTRY ABB=ON PLU=ON L25 NOT L24

=> d sta que 128

L1 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

VAR G5=H/AK

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 41

CONNECT IS M1 RC AT 45

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:


RSPEC 1

NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L2 2639 SEA FILE=REGISTRY CSS FUL L1

L21 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

```

VAR G5=H/AK
VAR G6=46/52/60
VAR G7=AK/CB/50/51
REP G8=(0-1) O
REP G9=(0-1) C
NODE ATTRIBUTES:
CONNECT IS M1 RC AT 45
CONNECT IS M1 RC AT 51
CONNECT IS M1 RC AT 57
CONNECT IS M1 RC AT 58
CONNECT IS M1 RC AT 59
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED
ECOUNT IS M1 O AT 57

```

```

GRAPH ATTRIBUTES:
RSPEC 1
NUMBER OF NODES IS 58

```

```


STEREO ATTRIBUTES: NONE
L23      324 SEA FILE=REGISTRY SUB=L2 CSS FUL L21
L24      12 SEA FILE=REGISTRY ABB=ON PLU=ON L23 AND NC>=2
L25      64 SEA FILE=REGISTRY ABB=ON PLU=ON L23 AND NR>=7
L26      52 SEA FILE=REGISTRY ABB=ON PLU=ON L25 NOT L24
L27      299 SEA FILE=REGISTRY ABB=ON PLU=ON L23 NOT ((D OR T)/ELS OR
          11C# OR 13C# OR 14C# OR C11# OR C13# OR C14# OR LABELED)
L28      235 SEA FILE=REGISTRY ABB=ON PLU=ON L27 NOT (L24 OR L25 OR L26)

```

```

=> d sta que 137
L1      STR

```



```

VAR G1=C/38/29
VAR G2=AK/OH/33/36
VAR G3=C/38
VAR G4=C/38/29/42/44
VAR G5=H/AK
NODE ATTRIBUTES:
CONNECT IS M1 RC AT 41
CONNECT IS M1 RC AT 45
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

```

GRAPH ATTRIBUTES:


RSPEC 1

NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L2 2639 SEA FILE=REGISTRY CSS FUL L1

L21 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

VAR G5=H/AK

VAR G6=46/52/60

VAR G7=AK/CB/50/51

REP G8=(0-1) O

REP G9=(0-1) C

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 45

CONNECT IS M1 RC AT 51

CONNECT IS M1 RC AT 57

CONNECT IS M1 RC AT 58

CONNECT IS M1 RC AT 59

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

ECOUNT IS M1 O AT 57

GRAPH ATTRIBUTES:


RSPEC 1

NUMBER OF NODES IS 58

STEREO ATTRIBUTES: NONE

L23 324 SEA FILE=REGISTRY SUB=L2 CSS FUL L21

L29 STR



```

VAR G1=C/38/29
VAR G2=AK/OH/33/36
VAR G3=C/38
VAR G4=C/38/29/42/44
VAR G5=H/AK
NODE ATTRIBUTES:
CONNECT IS M1 RC AT 41
CONNECT IS M1 RC AT 45
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

```

```


GRAPH ATTRIBUTES:
RSPEC 1
NUMBER OF NODES IS 42

```

```

STEREO ATTRIBUTES: NONE
L30 2505 SEA FILE=REGISTRY SUB=L2 CSS FUL L29
L31 STR

```



```

VAR G1=C/38/29
VAR G2=AK/OH/33/36
VAR G3=C/38
VAR G4=C/38/29/42/44

```

VAR G5=H/AK

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 45

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:


RSPEC 1

NUMBER OF NODES IS 42

STEREO ATTRIBUTES: NONE

L32 352 SEA FILE=REGISTRY SUB=L30 CSS FUL L31

L34 STR



VAR G1=C/38/29

VAR G2=AK/OH/33/36

VAR G3=C/38

VAR G4=C/38/29/42/44

VAR G5=H/AK

VAR G6=46/52/60

VAR G7=AK/CB/50/51

REP G8=(0-1) O

REP G9=(0-1) C

NODE ATTRIBUTES:

CONNECT IS M1 RC AT 45

CONNECT IS M1 RC AT 51

CONNECT IS M1 RC AT 57

CONNECT IS M1 RC AT 58

CONNECT IS M1 RC AT 59

DEFAULT MLEVEL IS ATOM

DEFAULT ECLEVEL IS LIMITED

ECOUNT IS M1 O AT 57

GRAPH ATTRIBUTES:

RSPEC 1

NUMBER OF NODES IS 58

STEREO ATTRIBUTES: NONE

L35 339 SEA FILE=REGISTRY SUB=L2 CSS FUL L34

L36 15 SEA FILE=REGISTRY ABB=ON PLU=ON L35 NOT (L32 OR L23)

L37 14 SEA FILE=REGISTRY ABB=ON PLU=ON L36 NOT 14C

```
=> fil hcaplus
FILE 'HCAPLUS' ENTERED AT 07:40:03 ON 13 DEC 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)
```

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

```
FILE COVERS 1907 - 13 Dec 2007 VOL 147 ISS 25
FILE LAST UPDATED: 12 Dec 2007 (20071212/ED)
```

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> => d 171 bib abs hitind hitstr retable tot
```

```
L71 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
```

```
AN 2004:370948 HCAPLUS Full-text
```

```
DN 140:375358
```

```
TI Stereospecific reduction of sapogen-3-ones
```

```
IN Gunning, Philip James; Tiffin, Peter David
```

```
PA Phytotech Limited, UK
```

```
SO PCT Int. Appl., 41 pp.
```

```
CODEN: PIXXD2
```

```
DT Patent
```

```
LA English
```

```
FAN.CNT 1
```

| | PATENT NO. | KIND | DATE | APPLICATION NO. | DATE |
|----|---------------|--|----------|-----------------|--------------|
| | ----- | ---- | ----- | ----- | ----- |
| PI | WO 2004037845 | A1 | 20040506 | WO 2003-GB1780 | 20030428 <-- |
| | W: | AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW | | | |
| | RW: | GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG | | | |
| | CA 2503899 | A1 | 20040506 | CA 2003-2503899 | 20030428 <-- |
| | AU 2003224308 | A1 | 20040513 | AU 2003-224308 | 20030428 <-- |
| | EP 1558627 | A1 | 20050803 | EP 2003-720733 | 20030428 <-- |
| | R: | AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK | | | |
| | BR 2003015746 | A | 20050906 | BR 2003-15746 | 20030428 <-- |
| | CN 1723218 | A | 20060118 | CN 2003-824744 | 20030428 <-- |
| | JP 2006507360 | T | 20060302 | JP 2005-501542 | 20030428 <-- |

| | | | | | |
|------|---|----|----------|----------------|--------------|
| | IN 2005MN00308 | A | 20060505 | IN 2005-MN308 | 20050420 <-- |
| | MX 2005PA04494 | A | 20050726 | MX 2005-PA4494 | 20050427 <-- |
| | US 2006041119 | A1 | 20060223 | US 2005-531086 | 20050621 <-- |
| | IN 2007MN01247 | A | 20071019 | IN 2007-MN1247 | 20070817 <-- |
| PRAI | GB 2002-25106 | A | 20021028 | <-- | |
| | GB 2003-1505 | A | 20030122 | | |
| | WO 2003-GB1780 | W | 20030428 | | |
| | IN 2005-MN308 | A3 | 20050420 | | |
| OS | CASREACT 140:375358; MARPAT 140:375358 | | | | |
| AB | A method to stereospecifically prepare a steroidal sapogenin or a derivative thereof by reducing a 3-keto,5 β -H steroidal sapogenin with a hindered organoborane or an organo-aluminum hydride. A 3 β -hydroxy,5 β -H steroidal sapogenin or derivative may be prepared by reducing the 3-keto,5 β -H steroidal sapogenin using as reducing agent which is a relatively highly hindered organoborane reagent or by SN 2 inversion of a 3 α -hydroxy,5 β -H steroidal sapogenin or derivative. The organo-aluminum hydride may be used to prepare a 3 α ,hydroxy,5 β -H steroidal sapogenin or derivative. The invention provides a convenient route to useful steroidal sapogenins such as sarsasapogenin, episarsasapogenin, smilagenin, epismilagenin and esters thereof, from readily available or easily prepared starting materials (e.g. diosgenone, prepared from diosgenin). | | | | |
| IC | ICM C07J0071-00 | | | | |
| CC | 32-8 (Steroids) | | | | |
| ST | stereospecific redn sapogenone; steroidal sapogenin prepn; smilagenin prepn; sarsasapogenin prepn | | | | |
| IT | Steroids, preparation | | | | |
| | RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation) | | | | |
| | (sapogenins; stereospecific reduction of sapogen-3-ones) | | | | |
| IT | Reduction | | | | |
| | (stereoselective; stereospecific reduction of sapogen-3-ones) | | | | |
| IT | Asymmetric synthesis and induction | | | | |
| | (stereospecific reduction of sapogen-3-ones) | | | | |
| IT | Sapogenins | | | | |
| | RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation) | | | | |
| | (steroidal; stereospecific reduction of sapogen-3-ones) | | | | |
| IT | 470-03-1P, Episarsasapogenin | | | | |
| | RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) | | | | |
| | (stereospecific reduction of sapogen-3-ones) | | | | |
| IT | 96-47-9, 2-Methyltetrahydrofuran 108-88-3, Toluene, uses 109-87-5, Dimethoxymethane 109-99-9, Thf, uses 123-91-1, 1,4-Dioxane, uses 1634-04-4, tert-Butyl methyl ether | | | | |
| | RL: NUU (Other use, unclassified); USES (Uses) | | | | |
| | (stereospecific reduction of sapogen-3-ones) | | | | |
| IT | 639-96-3, Sarsasapogenone 6870-79-7, Diosgenone | | | | |
| | RL: RCT (Reactant); RACT (Reactant or reagent) | | | | |
| | (stereospecific reduction of sapogen-3-ones) | | | | |
| IT | 126-19-2P, Sarsasapogenin 512-07-2P, Smilagenone 16653-88-6P, Epismilagenin | | | | |
| | RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent) | | | | |
| | (stereospecific reduction of sapogen-3-ones) | | | | |
| IT | 17476-04-9, Lithium tri-tert-butoxyaluminumhydride 38721-52-7, Lithium tri-sec-butylborohydride 54575-49-4, Potassium tri-sec-butylborohydride 60217-34-7, Lithium triamylborohydride 63717-74-8, Borate(1-), hydrotriphenyl-, | | | | |

lithium, (T-4)- 67276-04-4, Sodium tri-sec-butylborohydride
 67966-25-0, Potassium trisamylborohydride 99747-36-1,
 Potassium triphenylborohydride

RL: RGT (Reagent); RACT (Reactant or reagent)
 (stereospecific reduction of sapogen-3-ones)

IT 126-18-1P, Smilagenin 4952-69-6P,
 Smilagenin benzoate

RL: SPN (Synthetic preparation); PREP (Preparation)
 (stereospecific reduction of sapogen-3-ones)


IT 470-03-1P, Episarsasapogenin

RL: IMF (Industrial manufacture); RCT (Reactant); SPN (Synthetic
 preparation); PREP (Preparation); RACT (Reactant or reagent)
 (stereospecific reduction of sapogen-3-ones)

RN 470-03-1 HCAPLUS

CN Spirostan-3-ol, (3 α ,5 β ,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 639-96-3, Sarsasapogenone 6870-79-7, Diosgenone

RL: RCT (Reactant); RACT (Reactant or reagent)
 (stereospecific reduction of sapogen-3-ones)

RN 639-96-3 HCAPLUS

CN Spirostan-3-one, (5 β ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 6870-79-7 HCAPLUS


CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.




IT 126-19-2P, Sarsasapogenin 512-07-2P,
 Smilagenone 16653-88-6P, Epismilagenin
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (stereospecific reduction of sapogen-3-ones)
 RN 126-19-2 HCAPLUS
 CN Spirostan-3-ol, (3 β ,5 β ,25S)- (CA INDEX NAME)

Absolute stereochemistry.




RN 512-07-2 HCAPLUS
 CN Spirostan-3-one, (5 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 16653-88-6 HCAPLUS
 CN Spirostan-3-ol, (3 α ,5 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




IT 17476-04-9, Lithium tri-tert-butoxyaluminumhydride
 38721-52-7, Lithium tri-sec-butylborohydride 54575-49-4,
 Potassium tri-sec-butylborohydride 60217-34-7, Lithium
 triamylborohydride 63717-74-8, Borate(1-), hydrotriphenyl-,
 lithium, (T-4)- 67276-04-4, Sodium tri-sec-butylborohydride
 67966-25-0, Potassium trisamylborohydride 99747-36-1,
 Potassium triphenylborohydride
 RL: RGT (Reagent); RACT (Reactant or reagent)
 (stereospecific reduction of sapogen-3-ones)
 RN 17476-04-9 HCAPLUS
 CN Aluminate(1-), hydrotris(2-methyl-2-propanolato)-, lithium (1:1), (T-4)-
 (CA INDEX NAME)



RN 38721-52-7 HCAPLUS
 CN Borate(1-), hydrotris(1-methylpropyl)-, lithium (1:1), (T-4)- (CA INDEX NAME)




RN 54575-49-4 HCAPLUS
 CN Borate(1-), hydrotris(1-methylpropyl)-, potassium (1:1), (T-4)- (CA INDEX NAME)




RN 60217-34-7 HCAPLUS

CN Borate(1-), tris(1,2-dimethylpropyl)hydro-, lithium, (T-4)- (9CI) (CA INDEX NAME)




RN 63717-74-8 HCAPLUS

CN Borate(1-), hydrotriphenyl-, lithium, (T-4)- (9CI) (CA INDEX NAME)



RN 67276-04-4 HCAPLUS


CN Borate(1-), hydrotris(1-methylpropyl)-, sodium (1:1), (T-4)- (CA INDEX NAME)



RN 67966-25-0 HCAPLUS


CN Borate(1-), tris(1,2-dimethylpropyl)hydro-, potassium, (T-4)- (9CI) (CA

INDEX NAME)



RN 99747-36-1 HCAPLUS


CN Borate(1-), hydrotriphenyl-, potassium, (T-4)- (9CI) (CA INDEX NAME)

IT 126-18-1P, Smilagenin 4952-69-6P,
Smilagenin benzoateRL: SPN (Synthetic preparation); PREP (Preparation)
(stereospecific reduction of sapogen-3-ones)

RN 126-18-1 HCAPLUS

CN Spirostan-3-ol, (3β,5β,25R)- (CA INDEX NAME)


Absolute stereochemistry.




RN 4952-69-6 HCAPLUS

CN Spirostan-3-ol, benzoate, (3β,5β,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L71 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1983:405899 HCAPLUS Full-text
 DN 99:5899
 TI Modified steroids. Communication XII. Study of the Baeyer-Villiger reaction in a series of derivatives of a steroid compound diosgenin
 AU Irismetov, M. P.; Goryaev, M. I.; Rustambekova, G. B.; Mirzasaliev, N. A.
 CS Inst. Khim. Nauk, Alma-Ata, USSR
 SO Izvestiya Akademii Nauk Kazakhskoi SSR, Seriya Khimicheskaya (1983), (1), 75-7
 CODEN: IKAKAK; ISSN: 0002-3205
 DT Journal
 LA Russian
 GI



AB Diosgenones I underwent Baeyer-Villiger oxidation by BzO₂H in CHCl₃ to give lactones II, which were reduced by LiAlH₄ to give diols III.
 CC 32-8 (Steroids)
 ST diosgenone Baeyer Villiger oxidn; homooxaspirostanone lactone prepn redn; secospirostanediol
 IT Steroids, reactions
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (Baeyer-Villiger oxidation of diosgenins)
 IT Oxidation
 (Baeyer-Villiger, of diosgenins, homooxaspirostanones from)


IT 470-07-5 85881-65-8
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (Baeyer-Villiger oxidation of)
 IT 512-07-2P 85853-07-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation and reduction-ring cleavage of)
 IT 470-07-5
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (Baeyer-Villiger oxidation of)
 RN 470-07-5 HCAPLUS
 CN Spirostan-3-one, (5 α ,25R)- (CA INDEX NAME)

Absolute stereochemistry.



IT 512-07-2P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation and reduction-ring cleavage of)
 RN 512-07-2 HCAPLUS
 CN Spirostan-3-one, (5 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L71 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1960:74810 HCAPLUS Full-text
 DN 54:74810
 OREF 54:14301g-h
 TI A new method for the preparation of diosgenone
 AU Chakravarti, R. N.; Mitra, M. N.; Chakravarti, Debi
 CS Bethune Coll., Calcutta

- SO Bulletin of the Calcutta School of Tropical Medicine (1959), 7,
145
CODEN: BCSTA4; ISSN: 0068-5372
- DT Journal
- LA Unavailable
- OS CASREACT 54:74810
- AB Diosgenone (isospirost-4-en-3-one) was obtained from diosgenin by dissolving the latter (2 g.) in 40 ml. freshly distilled p-cymene and adding 1 g. Raney Ni in a 250- ml. flask fitted with an air condenser. The mixture was refluxed 12 hrs. in an atmosphere of dry N₂, was filtered hot, and the solvent removed by distillation under reduced pressure at 125-130°. The residue (1.4 g.) was chromatographed over Al₂O₃, and the crystalline solid eluted with 2:1 petr. ether-C₆H₆. The product was further purified by recrystn. from alc. The method was applicable to the preparation of 4-cholesten-3-one from cholesterol.
- CC 10J (Organic Chemistry: Steroids)
- IT Ultraviolet and visible, spectra
(of diosgenin)
- IT 601-57-0P, Cholest-4-en-3-one 6870-79-7P, Diosgenone
RL: PREP (Preparation)
(preparation of)
- IT 6870-79-7P, Diosgenone
RL: PREP (Preparation)
(preparation of)
- RN 6870-79-7 HCAPLUS
- CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.




- L71 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
- AN 1942:20583 HCAPLUS Full-text
- DN 36:20583
- OREF 36:3182i,3183a-c
- TI Sterols. CXXXIX. Sapogenins. 59. The bio-reduction of
4-dehydrotigogenone
- AU Marker, Russell E.; Wittbecker, Emerson L.; Wagner, R. B.; Turner, D. L.
- SO Journal of the American Chemical Society (1942), 64, 818-22
CODEN: JACSAT; ISSN: 0002-7863
- DT Journal
- LA Unavailable
- AB A 20-kg. male dog was fed daily (for 3 consecutive days) a mixture of 300 g. of dog biscuits and 30 g. of lard containing 3 g. of 4-dehydrotigogenone (I) and in addition 1 g. of I in 20 cc. peanut oil was injected subcutaneously; the feces were extracted with Me₂CO and ether and the residue from the extract was hydrolyzed with alc. KOH; the nonsaponifiable fraction (9.5 g.) gave 9.5 g. of a digitonin (II) precipitate, which yielded 0.2 g. of diosgenin and 0.1

g. of smilagenin; the fraction not precipitated by II contained 4.2 g. of unchanged I and 0.4 g. of epismilagenin, C₂₉H₄₆O₄ (III), m. 217-20° (acetate, m. 158-60°), separated as the succinic ester. III was also prepared from smilagenone by catalytic reduction (PtO₂ in EtOH for 75 min. at room temperature) or by the action of Na in absolute EtOH. III is reoxidized to IV by CrO₃ in 90% AcOH. Further reduction (PtO₂ in AcOH at 70-5° and 3 atmospheric of H for 10 hrs.) gives epidihydrosarsapogenin, m. 134-6°; crystallization from Me₂CO gives a polymorphic form, m. 180-2°. The dog normally excretes epicoprosterol in a considerable amount; this lends addnl. support to Schoenheimer's theory (C. A. 32, 7985.2) that cholestenone is an intermediate in the formation of coprosterol in the organism. The significance of these facts is discussed.


CC 10 (Organic Chemistry)
 IT Animal organism
 (4-dehydrotigogenone reduction in)
 IT Reduction
 (of 4-dehydrotigogenone in animal organism)
 IT Sarsasapogenin, epidihydro-
 RL: PREP (Preparation)
 IT 6870-79-7, Tigogenone, 4-dehydro-
 (bio-reduction of)
 IT 126-18-1P, Smilagenin 512-04-9P,
 Diosgenin 512-07-2P, Smilagenone 16653-88-6P,
 Epismilagenin 106759-14-2P, Epismilagenin,
 acetate
 RL: PREP (Preparation)
 (preparation of)
 IT 6870-79-7, Tigogenone, 4-dehydro-
 (bio-reduction of)
 RN 6870-79-7 HCAPLUS
 CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.



IT 126-18-1P, Smilagenin 512-04-9P,
 Diosgenin 512-07-2P, Smilagenone 16653-88-6P,
 Epismilagenin 106759-14-2P, Epismilagenin,
 acetate
 RL: PREP (Preparation)
 (preparation of)
 RN 126-18-1 HCAPLUS
 CN Spirostan-3-ol, (3 β , 5 β , 25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 512-04-9 HCAPLUS

CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 512-07-2 HCAPLUS

CN Spirostan-3-one, (5 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 16653-88-6 HCAPLUS

CN Spirostan-3-ol, (3 α ,5 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 106759-14-2 HCAPLUS

CN Spirostan-3-ol, 3-acetate, (3 α ,5 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.



=> => d 195 all hitstr retable

L95 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1983:215889 HCAPLUS Full-text

DN 98:215889

OREF 98:32841a,32844a

ED Entered STN: 12 May 1984

TI Modified steroids. XI. Preparation of epoxy compounds from diosgenin and its derivatives

AU Irismetov, M. P.; Goryaev, M. I.; Rustembekova, G. B.; Mirzasaliev, N. A.

CS Inst. Khim. Nauk, Alma-Ata, USSR

SO Zhurnal Obshchei Khimii (1983), 53(2), 462-5


CODEN: ZOKHA4; ISSN: 0044-460X

DT Journal

LA Russian

CC 32-8 (Steroids)

GI



AB Epoxidn. of diosgenins I (R = H, Ac) and II (Z = O, H₂) by BzOOH in CHCl₃ gave epoxides III and IV, resp. LiAlH₄ reduction of III (R = H, Ac) and IV (Z = O) gave 5 α -spirostane-3 β ,5-diol.

ST epoxidn diosgenin; epoxyhydroxyspirostane

IT Steroids, reactions

RL: RCT (Reactant); RACT (Reactant or reagent)
(epoxidn. of diosgenins)

IT Epoxidation

(of diosgenins)

IT 512-04-9 6870-79-7 85707-30-8

RL: RCT (Reactant); RACT (Reactant or reagent)
(epoxidn. of)

IT 85707-31-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
(Reactant or reagent)
(preparation and acetylation of)

IT 1061-54-7P 3514-60-1P 66965-00-2P 85719-33-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP
(Preparation); RACT (Reactant or reagent)
(preparation and epoxide ring cleavage of)

IT 85707-32-0P 85707-33-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)


IT 512-04-9 6870-79-7

RL: RCT (Reactant); RACT (Reactant or reagent)
(epoxidn. of)

RN 512-04-9 HCAPLUS


CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.




RN 6870-79-7 HCAPLUS
 CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.



IT 1061-54-7P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP
 (Preparation); RACT (Reactant or reagent)
 (preparation and epoxide ring cleavage of)
 RN 1061-54-7 HCAPLUS
 CN Spirost-5-en-3-ol, acetate, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.



=> => d bib abs hitstr retable tot 197

L97 ANSWER 1 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1981:407585 HCAPLUS Full-text
 DN 95:7585
 OREF 95:1443a,1446a
 TI Study of synthetic transformations of a steroidal compound of diosgenin
 AU Irismetov, M. P.; Goryaev, M. I.
 CS USSR
 SO Trudy Instituta Khimicheskikh Nauk, Akademiya Nauk Kazakhskoi SSR (
 1980), 52, 17-39
 CODEN: TIKNAG; ISSN: 0568-5087
 DT Journal
 LA Russian
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Heterocyclic analogs of diosgenin were prepared. Thus, cyclocondensation of 2-formyldiosgenone with N_2H_4 and HONH_2 gave the pyrazalodiosgenin I and isoxazolodiosgenin II, resp. Fisher indole synthesis of diosgenone with PhNHNH_2 gave indolodiosgenin III, and Beckmann rearrangement of dihydrodiosgenone oxime gave lactams IV and V ($\text{Z} = \text{O}$), which were reduced by LiAlH_4 to give IV and V ($\text{Z} = \text{H}_2$). Baeyer-Villiger oxidation of dihydrodiosgenone gave lactam VI, and cyclocondensation of 2α -bromodihydrodiosgenone with PhCH:NNHC(S)NH_2 gave the thiazolodiosgenin VII.

IT 470-07-5P 512-07-2P


RL: RCT (Reactant); PREP (Preparation); RACT (Reactant or reagent)

(Fisher indole synthesis of)

RN 470-07-5 HCAPLUS

CN Spirostan-3-one, ($5\alpha, 25\text{R}$)- (CA INDEX NAME)


Absolute stereochemistry.



RN 512-07-2 HCAPLUS

CN Spirostan-3-one, ($5\beta, 25\text{R}$)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 512-04-9

RL: RCT (Reactant); RACT (Reactant or reagent)
(oxidation of)

RN 512-04-9 HCAPLUS


CN Spirost-5-en-3-ol, ($3\beta, 25\text{R}$)- (CA INDEX NAME)

Absolute stereochemistry.



IT 6870-79-7P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP
 (Preparation); RACT (Reactant or reagent)
 (preparation and acylation of)
 RN 6870-79-7 HCAPLUS
 CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.




L97 ANSWER 2 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1966:68078 HCAPLUS Full-text
 DN 64:68078
 OREF 64:12755e-h,12756a-b
 TI Structure of yononin. A novel type of spirostanol glycoside
 AU Kawasaki, T.; Miyahara, K.
 CS Kyushu Univ., Fukuoka, Japan
 SO Tetrahedron (1965), 21(12), 3633-9
 CODEN: TETRAB; ISSN: 0040-4020
 DT Journal
 LA English
 GI For diagram(s), see printed CA Issue.
 AB cf. CA 59, 15535g. Isolation from the rhizome of Dioscorea tokoro gave yononin (I), and α -L-arabinoside of yonogenin (II) (Takada, et al., CA 53, 16206f). I (200 mg.) in 5 ml. HCONMe₂ methylated with 2.3 g. MeI and 1 g. Ag₂O, the procedure repeated twice, and the sirup refluxed 2 hrs. with 4 ml. 2N HCl in 50% aqueous alc. gave 24 mg. aglycon (II), m. 184-6°, [α]_D -58° (c 0.55, all in CHCl₃), R_f 0.34 (3:1 C₆H₁₂-EtOAc, 10% H₂SO₄ spray). II must be a yonogenin monomethyl ether and synthetic 2- and 3-methyl ethers were prepared for comparison. Preparation from diosgenin gave 25D-spirost-4-en-3-one (III), m. 185-6° [α]_D 4° (c 0.52). III (5 g.) in 350 ml. MeOH treated 10 min. with 20 ml. 30% H₂O₂ and 20 ml. 4N NaOH, the mixture stirred 7 hrs. at 2° and kept 16 hrs. at 0° gave 4,5-epoxy-25D-spirostan-3-one, m. 205-7°, [α]_D 23° (c

0.53). The epoxide (2.98 g.) in 100 ml. Me₂CO treated dropwise with 6 ml. 25% H₂SO₄ and the mixture kept 4 days at 20° gave the 2 α -hydroxy-4-en-3-one (IV), m. 213.5-15.5° [α]_D 21° (c 0.65), methylated (1 g.) in 25 ml. dry C₆H₆ by stirring 20 hrs. with 3 g. MeI and 3 g. Ag₂O to give the 2 α -methoxy-4-en-3-one (V), m. 198-201°, [α]_D 29° (c 0.44). V (200 mg.) in 40 ml. alc. hydrogenated over 50 mg. 10% Pd-C gave 19 mg. 2 α -methoxy-25D-spirost-4-en-3-ol, m. 201-2°, [α]_D 11° (c 0.44), and 2 β -methoxy-25D,5 β -spirostan-3-one (VI), m. 212°. VI (20 mg.) reduced 4 hrs. by stirring in 2 ml. dry C₅H₅N containing 5 mg. LiBH₄ gave 2 β -methoxy-25D, 5 β -spirostan-3 β -ol, m. 205-8°, [α]_D -49° (c 0.51), and 2 β -methoxy-25D,5 β -spirostan-3 α -ol, m. 265-6°, [α]_D -119° (c 0.37), identical with II 2-methyl ether. III (770 mg.) acetylated overnight at 20° with 20 ml. 1:3 Ac₂O-C₅H₅N gave 740 mg. 2 α -acetoxy-4-en-3-one (VII), m. 248-50.5°, [α]_D -15° (c 0.66). VII (400 mg.) in 30 ml. EtOAc hydrogenated over 50 mg. 10% Pd-C 10 min. and the crystalline mass (395 mg.) reduced in 50 ml. MeOH with 100 mg. NaBH₄, the product (374 mg.) methylated in 15 ml. dry C₆H₆ with 300 mg. MeI and 500 mg. Ag₂O for 40 hrs., and the resultant Me ether acetate refluxed 40 min. with 30 ml. 5% KOH-MeOH gave a mixture of 3-methoxy-25D-spirostan-2-ols. The mixture (340 mg.) chromatographed on 10 g. Al₂O₃ and eluted with C₆H₆ gave 248 mg. 3 α -methoxy-25D,5 β -spirostan-2 α -ol (VIII), m. 162-3°, [α]_D -31° (c 0.70). VIII (16 mg.) in 1 ml. 90% AcOH oxidized with 0.1 ml. solution (200 mg. CrO₃ in 1 ml. 90% AcOH) under stirring 4.5 hrs. gave 10 mg. 3 α -methoxy-25D,5 β -spirostan-2-one (IX), m. 198°, [α]₃₁₄ -741°, [α]₂₈₀ 36° (c 0.305, MeOH). IX (50 mg.) in 5 ml. MeOH treated with NaBH₄ gave 42 mg. compound, m. 161 .apprx. 2°. IX (71 mg.) reduced with LiBH₄ in C₅H₅N gave VIII and 3 α -methoxy-25D,5 β -spirostan-2 β -ol, m. 185 .apprx. 8°, [α]_D -58° (c 0.63), identical with II. Consequently, I is defined as O- α -L-arabinosyl- (1-2)-25D,5 β -spirostan-2 β ,3 α -diol (yonogenin 2- α -L-arabinoside). This is the 1st spirostanol glycoside shown to have the sugar moiety combined with an OH group other than that at C-3 of the aglycon.

IT 5247-71-2P, 5 β ,25D-Spirostan-2 β -ol, 3 α -methoxy-
 5247-73-4P, 25D-Spirost-4-en-3-one, 2 α -hydroxy-
 5247-75-6P, 5 β ,25D-Spirostan-2 α -ol, 3 α -methoxy-
 5289-76-9P, 5 β ,25D-Spirostan-3 α -ol, 2 β -methoxy-
 5372-57-6P, 25D-Spirost-4-en-3-one, 2 α -methoxy-
 5373-18-2P, 5 β ,25D-Spirostan-3-one, 2 β -methoxy-
 5605-39-0P, 25D-Spirost-4-en-3-ol, 2 α -methoxy-
 5605-40-3P, 5 β ,25D-Spirostan-3 β -ol, 2 β -methoxy-
 6870-79-7P, 25D-Spirost-4-en-3-one
 RL: PREP (Preparation)
 (preparation of)
 RN 5247-71-2 HCAPLUS
 CN Spirostan-2-ol, 3-methoxy-, (2 β ,3 α ,5 β ,25R)- (9CI) (CA
 INDEX NAME)


Absolute stereochemistry.



RN 5247-73-4 HCAPLUS

CN 25D-Spirost-4-en-3-one, 2 α -hydroxy-, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5247-75-6 HCAPLUS

CN 5 β ,25D-Spirostan-2 α -ol, 3 α -methoxy-, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5289-76-9 HCAPLUS

CN 5 β ,25D-Spirostan-3 α -ol, 2 β -methoxy-, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5372-57-6 HCAPLUS

CN Spirost-4-en-3-one, 2-methoxy-, (2 α ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5373-18-2 HCAPLUS

CN 5 β ,25D-Spirostan-3-one, 2 β -methoxy-, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5605-39-0 HCAPLUS

CN 25D-Spirost-4-en-3-ol, 2 α -methoxy-, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5605-40-3 HCAPLUS

CN Spirostan-3-ol, 2-methoxy-, (2 β , 3 β , 5 β , 25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 6870-79-7 HCAPLUS

CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 3 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1963:428741 HCAPLUS Full-text

DN 59:28741

OREF 59:5234b-g

TI Steroidal components of domestic plants. XL. Constituents of *Heloniopsis orientalis*. 3. The structure of heloniogenin

AU Okanishi, Tameto; Akahori, Akira; Yasuda, Fumio

CS Shionogi Co., Osaka, Japan

SO Chemical & Pharmaceutical Bulletin (1962), 10, 1195-9

CODEN: CPBTAL; ISSN: 0009-2363

DT Journal

LA Unavailable

GI For diagram(s), see printed CA Issue.


AB cf. Shionogi Kenkyusho Nempo 11, 97-101(1961); CA 57,8637h. In addition to the properties and derivs. of heloniogenin (I) previously reported [ibid. 10, 1411-15(1960)], its monoacetylation and CrO₃ oxidation were described. I (500 mg.) kept with 4 ml. Ac₂O in C₅H₅N 2 hrs. at 10° and the mixture poured into ice H₂O, and extracted with ether yielded 572 mg. residue from the extract, which was separated by Al₂O₃ chromatography into 10 mg. previously reported diacetate, m. 184-5°, 180 mg. unchanged I, m. 212-13°, and 225 mg. desired 3-acetate (II) of I, m. 218-19°, [α]₂₅ -89.6 \pm 2° (c 1.029, CHCl₃). Oxidation of 280 mg. I with CrO₃ in AcOH 30 min. at room temperature yielded 261 mg. mixture, which was separated by Al₂O₃ chromatography into 21 mg. 25D-spirost-4-ene-3,12-dione (III), m. 248-50°, and 42 mg. gentrogenin (3 β -hydroxy-25D-spirost-5-en-12-one) (IV), m. 215-16°, [α]_{28D} -56.0 \pm 2° (c 1.021, CHCl₃), and 42 mg. mixture of I and IV. Similar oxidation of 450 mg. II yielded 420 mg. mixture, which was separated by Al₂O₃ chromatography into 240 mg. IV acetate, m. 224-5°, [α]_{25D} -58.1 \pm 2° (c 1.016, CHCl₃), and 150 mg. recovered II. These results showed I to be 25D-spirost-5-ene-3 β ,12 ξ -diol. The configuration of the 12-OH group remained to be determined. IV acetate (300 mg.) reduced with LiAlH₄ in ether in the usual way yielded 317 mg. mixture of isomers, separated by Al₂O₃ chromatography into 165 mg. 3 β ,12 β -diol (V) [acetate (VI) m. 206-7°, [α]_{27D} -117.8 \pm 2° (c 1.015, CHCl₃)] and 105 mg. 3 β ,12 α -diol (VII), m. 211-12°, [α]_{23D} -89.2 \pm 2° (c 1.062, CHCl₃); diacetate (VIII) m. 180-2°, [α]_{27D} -61.8 \pm 2° (c 1.080, CHCl₃). VI (50 mg.) hydrolyzed with KOH-EtOH yielded 45 mg. V, m. 233-5°, [α]_{27D} -116.4 \pm 2° (c 0.993, CHCl₃). IV acetate (230 mg.) reduced with NaBH₄ in EtOH gave similar results and yielded 130 mg. V and 50 mg. VII. V and VI showed the same phys. consts. as isochiapagenin (IX) (m. 236-7°, [α]_D -121°) and its acetate (m. 206-7°, [α]_D -120°), obtained from chiapagenin by refluxing 94 hrs. with HCl EtOH, and acetylating the resulting IX with Ac₂O-C₅H₅N. VII and VIII were identical with I and I diacetate, resp., in m.p., [α]_D, and infrared spectra, and showed no depression of mixed m.p. The structure of I was thus established as 25D-spirost-5-ene-3 β ,12 α -diol.

IT 6869-16-5, 25D-Spirost-5-ene-3 β ,12 α -diol
(as structure for heloniogenin)

RN 6869-16-5 HCAPLUS

CN Spirost-5-ene-3,12-diol, (3 β ,12 α ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 427-28-1P, Gentrogenin 5996-01-0P, Gentrogenin, acetate
6875-60-1P, 25D-Spirost-4-ene-3,12-dione 6877-71-0P,
25D-Spirost-5-ene-3 β ,12 β -diol 59203-51-9P,


Isochiapagenin 103592-10-5P, Heloniogenin, 3-acetate
 105063-79-4P, Heloniogenin, diacetate 105859-99-2P,
 25D-Spirost-5-ene-3 β ,12 β -diol, diacetate
 RL: PREP (Preparation)

(preparation of)

RN 427-28-1 HCAPLUS

CN Spirost-5-en-12-one, 3-hydroxy-, (3 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 5996-01-0 HCAPLUS

CN Spirost-5-en-12-one, 3-(acetyloxy)-, (3 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 6875-60-1 HCAPLUS

CN Spirost-4-ene-3,12-dione, (25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 6877-71-0 HCAPLUS

CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 59203-51-9 HCAPLUS

CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,20 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 103592-10-5 HCAPLUS

CN Heloniogenin, 3-acetate (7CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 105063-79-4 HCAPLUS

CN Spirost-5-ene-3,12-diol, diacetate, (3 β ,12 α ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 105859-99-2 HCAPLUS

CN 25D-Spirost-5-ene-3 β ,12 β -diol, diacetate (7CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 6869-16-5, Heloniogenin
(structure of)

RN 6869-16-5 HCAPLUS

CN Spirost-5-ene-3,12-diol, (3 β ,12 α ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 4 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1962:443009 HCAPLUS Full-text

DN 57:43009

OREF 57:8637h-i,8638a-h

TI Steroidal components of domestic plants. XXXII. Constituents of Reineckia carnea. 4. Structure of kitigenin. 1

AU Sasaki, Kanzo

CS Shionogi & Co., Osaka

SO Chemical & Pharmaceutical Bulletin (1961), 9, 684-92

CODEN: CPBTAL; ISSN: 0009-2363

DT Journal

LA Unavailable

OS CASREACT 57:43009

GI For diagram(s), see printed CA Issue.

AB cf. CA 56, 11664a. (Throughout these abstrs. optical rotations were measured in CHCl_3). The positions and configurations of 3 of the 4 OH groups of kitigenin (I), previously shown (loc. cit.) to be in ring A, were determined. Acetylation of 588 mg. I with Ac_2O - $\text{C}_5\text{H}_5\text{N}$ yielded 724 mg. mixture, separated by CS_2 into 370 mg. CS_2 -insol. diacetate (II), m. $217-19^\circ$, $[\alpha]_{29.5\text{D}} -45.6 \pm 2^\circ$ (c 0.868), and 354 mg. CS_2 -soluble triacetate (III), m. $219-20.5^\circ$ (depressed when mixed with II), $[\alpha]_{30\text{D}} -53.6 \pm 2^\circ$ (c 1.023). II (0.2 g.) in Me_2CO treated dropwise with CrO_3 - H_2SO_4 yielded 225 mg. ketone (IV), m. $200-2^\circ$, $[\alpha]_{24\text{D}} -64.7 \pm 2^\circ$ (c 1.002). Thus, only 1 OH group in II was oxidized, as evidenced by both ultraviolet and infrared spectra, and the remaining OH group must be tertiary, at C-5. III in $\text{C}_5\text{H}_5\text{N}$ was dehydrated with SOCl_2 to give 74% the 4-ene triacetate (V), m. $218-21^\circ$ (depressed when mixed with III), $[\alpha]_{24\text{D}} -129 \pm 2^\circ$ (c 1.105). Infrared bands at 1762 and 1730 cm^{-1} confirmed the presence of the enol acetate and acetoxy groups, resp., thus proving an AcO group at C-4. Reductive hydrolysis of 0.1 g. V in tetrahydrofuran with LiAlH_4 in ether yielded 81 mg. oxo diol (VI), m. $192-6^\circ$, which gave a pos. ketol test with 2,3,5-triphenyltetrazolium chloride, and a neg. FeCl_3 color test. VI (48 mg.) oxidized with $(\text{AcO})_2\text{Cu}$ in AcOH - MeOH yielded 45 mg. 4-hydroxy-1,4-dien-3-one compound (VII), m. $175-95^\circ$, which gave a neg. ketol test and a pos. FeCl_3 color test, and was acetylated with Ac_2O - $\text{C}_5\text{H}_5\text{N}$ to yield 15 mg. 4-acetoxy-1,4-dien-3-one compound (VIII), m. $212-15^\circ$. VII (46 mg.) was also formed by refluxing 62 mg. VI with 3% NaOH - MeOH . Catalytic reduction ($\text{Pd}-\text{C}$) of VII in AcOEt gave 4-hydroxy-25D-spirost-4-en-3-one (IX), m. $215-19^\circ$, $[\alpha]_{29\text{D}} 10.3 \pm 6^\circ$ (c 0.361). The structure of IX was confirmed by its synthesis from 0.4 g. diosgenone (X) by oxidation with H_2O_2 in the presence of OsO_4 in ether, whereby 287 mg. X was recovered and 20 mg. IX obtained, identical with the preceding sample. The yield of IX was increased to 30% by increasing the reaction time from 24 to 90 hrs. These results indicated the location of 3 of the 4 OH groups at the 3-, 4-, and 5-positions in I. The configuration of the OH group at C-3 was next determined. II (0.2 g.) in $\text{C}_5\text{H}_5\text{N}$ kept overnight at 0° with MeSO_2Cl yielded 98 mg. unsatd. triol diacetate (XI), m. $236-40^\circ$, which (22 mg.) was hydrolyzed with 1.5% NaOH - MeOH to yield 20 mg. unsatd. triol (XII), m. $235-9^\circ$, $[\alpha]_{24\text{D}} 18.7 \pm 4^\circ$ (c 0.503). XI (0.2 g.) catalytically hydrogenated (PtO_2) in AcOH yielded 196 mg. mixture, separated by Al_2O_3 chromatography into a trace of (probably) the 4,5-diol monoacetate, m. $195-212^\circ$, and 140 mg. corresponding saturated triol diacetate (XIII), m. $204-6^\circ$, $[\alpha]_{23\text{D}} -21.1 \pm 2^\circ$ (c 0.848). X (5 g.) reduced with LiAlH_4 in ether yielded 5.02 crude epimeric mixture of 4-en-3-ols, which (1.08 g.) was separated by digitonin precipitation into the 4-en-3 α -ol (XIV), m. $182-4^\circ$, $[\alpha]_{20\text{D}} -5.1 \pm 3^\circ$ (c 0.831), and the 4-en-3 β -ol (XV), m. $155-7^\circ$, $[\alpha]_{23\text{D}} -39.6 \pm 3^\circ$ (c 0.727) in a 2.5:1 ratio. Acetylation of XIV and XV with Ac_2O - $\text{C}_5\text{H}_5\text{N}$ gave the corresponding 4-en-3 α -acetate (XVI), m. $170-2^\circ$, $[\alpha]_{18\text{D}} 89.3 \pm 2^\circ$ (c 1.035), and the 4-en-3 β -acetate (XVII), m. $167-9^\circ$, $[\alpha]_{22\text{D}} -82.0 \pm 2^\circ$ (c 1.019). The $[\text{M}]_D$ differences between the alcs. and their acetates, when compared with those of cholest-4-en-3-ols and their acetates, supported the assigned configurations. cis-Hydroxylation of 1.01 g. XV with OsO_4 yielded 0.86 g. mixture separated into 508 mg. C_6H_6 -soluble fraction (A) and 376 mg. C_6H_6 -insol. fraction (B). Chromatography of A over Florisil gave 332 mg. recovered XV and 97 mg. triol mixture, which with B was acetylated with Ac_2O - $\text{C}_5\text{H}_5\text{N}$ to yield 540 mg. acetate mixture, and this chromatographed over Al_2O_3 yielded 27 mg. 3 β ,4 β ,5 β -triol 3,4-diacetate, m. $200-3^\circ$ (identical by mixed m.p. and infrared spectra with XIII obtained indirectly from I), and 425 mg. 3 β ,4 α ,5 α -triol 3,4-diacetate (XVIII), m. $245.5-6.0^\circ$, $[\alpha]_{23\text{D}} -23.4 \pm 2^\circ$ (c 0.603).

Infrared data were reported in support of the structures of II-IX and XI-XVIII. Thus was established the β -orientation of the 3-OH group, and the cis-relation of the 4- and 5-OH groups.


IT 104759-91-3 107297-14-3 107656-53-1
107741-57-1

(Derived from data in the 7th Collective Formula Index (1962-1966))

RN 104759-91-3 HCAPLUS

CN 25D-Spirost-4-en-3-ol, acetate (7CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 107297-14-3 HCAPLUS

CN 5 α , 25D-Spirostan-1 α , 3 α -diol (7CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 107656-53-1 HCAPLUS


CN 5 α , 25D-Spirostan-1 α , 3-diol, 3-acetate (7CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 107741-57-1 HCAPLUS
 CN 25D-Spirost-4-en-3-ol (7CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 13944-32-6P, 25D-Spirost-4-en-3-one, 4-hydroxy-
 16653-41-1P, 25D-Spirost-4-en-3 β -ol 16653-54-6P,
 25D-Spirost-4-en-3 β -ol, acetate 106505-71-9P,
 25D-Spirosta-1,4-dien-3-one, 4-hydroxy-
 RL: PREP (Preparation)
 (preparation of)
 RN 13944-32-6 HCAPLUS
 CN Spirost-4-en-3-one, 4-hydroxy-, (25R)- (8CI, 9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 16653-41-1 HCAPLUS
 CN Spirost-4-en-3-ol, (3 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 16653-54-6 HCAPLUS

CN Spirost-4-en-3 β -ol, acetate, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 106505-71-9 HCAPLUS

CN 25D-Spirosta-1,4-dien-3-one, 4-hydroxy- (7CI) (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 5 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1961:106030 HCAPLUS [Full-text](#)

DN 55:106030

OREF 55:19989b-i,19990a-b

TI Chiapagenin and isochiapagenin. Two new steroidal sapogenins from *Dioscorea chiapasensis*

AU Harrison, I. T.; Velasco, M.; Djerassi, Carl

CS Stanford Univ., Stanford, CA

SO Journal of Organic Chemistry (1961), 26, 155-8

CODEN: JOCEAH; ISSN: 0022-3263

DT Journal

LA Unavailable

AB Two new dihydroxy sapogenins, chiapagenin (I) and isochiapagenin (II), isolated from *D. chiapasensis*, were shown by appropriate interconversions to be 12 β -hydroxyyamogenin and 12 β -hydroxydiosgenin, resp. Dried and powdered roots (1 kg.) refluxed 2 hrs. in 10 l. denatured alc., the extraction repeated twice, and the combined exts. concentrated to 5 l., refluxed 4 hrs. with 1.5 l. concentrated HCl, diluted with 30 l. ice H₂O and the washed precipitate dried in vacuo at 80° gave 52 g. sapogenins. The product (25 g.) and 4.0 g. Girard reagent T refluxed 1 hr. in 165 ml. 10:1 alc.-AcOH, the cooled solution added to excess saturated aqueous NaHCO₃ and unchanged sapogenins (III) (23.5

g.) removed by 3-fold extraction with Et₂O, the aqueous layer adjusted to pH 1.0 with concentrated HCl and heated 1 hr. on a steam bath, the cooled mixture extracted with Et₂O and the residue on evaporation chromatographed on 60 g. Al₂O₃ (activity III), eluted with C₆H₆ and the fraction crystallized from alc., acetylated, and recrystd. from Et₂O-C₆H₁₄ gave 13 mg. correlogenin (IV) acetate, m. 211-12°. III (8.8 g.) chromatographed on 350 g. Al₂O₃, eluted with 500 ml. C₆H₆ and the fraction crystallized from alc. gave 0.46 g. I 3,5-diene derivative, m. 194-5°, [α]_D -188° (c 1.1), λ 227, 235, 242 mμ, produced by dehydration during the acid hydrolysis. Further elution with 8:2 C₆H₆-Et₂O gave 1.78 g. yamogenin (V) and diosgenin (VI) mixture, m. 180-7°. The acetylated mixture (2.1 g.) chromatographed on 80 g. Al₂O₃ (activity II) and eluted with 1:1 C₆H₆-C₆H₁₄ gave 110 mg. material, m. 179-84°, rechromatographed and recrystd. to yield 21 mg. pure VI acetate, m. 196-7°, [α]_D -126°, and 370 mg. material, m. 174-6°, recrystd. from alc. to give 219 mg. V acetate, m. 177-8°, [α]_D -126°, hydrolyzed to V, m. 195-6°. Further elution of the column with 6:4 C₆H₆-Et₂O gave 4.03 g. fraction, recrystd. from alc. to yield 3.8 g. material, m. 204-5°, converted to the diacetate, m. 194-6° (alc.), [α]_D -128° (c 1.9), hydrolyzed to give pure I, m. 257-9°, [α]_D -130° (c 1.2), ν 922, 895, 853 cm.⁻¹ (CS₂), indicative of the neo rather than the iso side chain configuration. I (450 mg.) and 3.1 ml. cyclohexanone in 20 ml. PhMe distilled with passage of some solvent and the mixture refluxed gently 4 hrs. with addition of 315 mg. Al(OCHMe₂)₃ in 2 ml. PhMe, diluted with H₂O and extracted with Et₂O gave 190 mg. ketone (VII), m. 214-17° (C₆H₆-C₆H₁₄), [α]_D -13° (c 1.0), λ 240 mμ (ε 16,800, alc.). I (480 mg.) kept 2 hrs. at 20° in 4 ml. Ac₂O and 25 ml. C₅H₅N, the monoacetate taken up in 5 ml. Et₂O, filtered and the Et₂O-soluble fraction chromatographed on 20 g. Al₂O₃ (activity I), eluted with 4:1 C₆H₆-Et₂O and recrystd. from aqueous MeOH yielded 325 mg. I 3-monoacetate (VIII), m. 176-7°, [α]_D -119° (c 0.4). VIII (213 mg.) in 10 ml. AcOH at 10° kept 30 min. with 53 mg. CrO₃ in 25 ml. AcOH, diluted with H₂O and Et₂O, the washed and dried organic phase evaporated and the residue crystallized from dilute MeOH gave 151 mg. IV acetate. I diacetate (2.17 g.) in 50 ml. AcOH hydrogenated 2 hrs. with 100 mg. prereduced PtO₂ and the reduction product chromatographed on 100 g. Al₂O₃ gave dihydrochiapagenin diacetate (IX), m. 204-5°, [α]_D -76° (c 0.6), saponified with boiling 5% KOH in MeOH to dihydrochiapagenin (X), m. 202-4° (dilute MeOH), [α]_D -79° (c 1.1). IX (33 mg.) and 11.5 g. LiOH.H₂O in 40 ml. 80% alc. kept 22 hrs. at 21°, diluted with H₂O and extracted with Et₂O gave dihydrochiapagenin 12-monoacetate (XI), m. 213-14° (C₆H₆-C₆H₁₄), [α]_D -84° (c 0.3). Sisalagenin acetate (109 mg.) refluxed 2 hrs. in absolute alc. with 13 mg. NaBH₄, the mixture refluxed 1 hr. with 100 mg. NaOH, diluted with H₂O and extracted with Et₂O gave X, m. 204-5° (dilute MeOH), [α]_D -73° (c 0.7), m. 194-6° (polymorphic form), acetylated and recrystd. from MeOH to give IX. Selective saponification of IX with LiOH.H₂O gave XI. A new and larger batch of *D. chiapasensis* (7 kg.) was extracted to yield 67 g. crystalline and 63 g. oily sapogenin mixture. Chromatography of the crystalline fraction gave 13 g. V-VI mixture and 22 g. I. Similar chromatography of the oily fraction produced 20 g. I 3,5-diene derivative, 14 g. V-VI mixture and, after acetylation, 0.535 g. II 3,12-diacetate, m. 206-7° (C₆H₁₄), [α]_D -120° (c 1.3), saponified in boiling 5% alc. NaOH and recrystd. from MeOH to yield II, m. 236-7°, [α]_D -121° (c 0.8), monoacetylated (101 mg.) to yield 60 mg. II 3-monoacetate, m. 208-10° (C₆H₁₄). Oxidation of 50 mg. monoacetate with 15 mg. CrO₃ gave 31 mg. botogenin acetate, m. 226-7°, [α]_D -56° (c 0.88), reduced (97 mg.) in 10 ml. absolute alc. with 13 mg. NaBH₄ by refluxing 2 hrs. and acetylated to yield II acetate, m. 206-7°.


II 1180-12-7 118923-52-7 119008-69-4
119719-99-2 120576-49-0 121009-56-1
125590-14-9

(Derived from data in the 6th Collective Formula Index (1957-1961))

RN 1180-12-7 HCAPLUS

CN Spirost-5-en-3-ol, acetate, (3 β ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 118923-52-7 HCAPLUS

CN 25L-Spirost-5-ene-3 β ,12 β -diol, diacetate (6CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 119008-69-4 HCAPLUS

CN 5 α ,25L-Spirostan-3 β ,12 β -diol, diacetate (6CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 119719-99-2 HCAPLUS

CN Isochiapagenin, 3-acetate (6CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 120576-49-0 HCAPLUS

CN Spirostan-3,12-diol, 12-acetate, (3 β ,12 β ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 121009-56-1 HCAPLUS

CN 5 α ,25L-Spirostan-3 β ,12 β -diol, 12-acetate (6CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 125590-14-9 HCAPLUS


CN Chiapagenin, 3-acetate (6CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 512-06-1P, Yamogenin
 RL: PREP (Preparation)
 (acetate and separation of yamogenin, from *Dioscorea chiapasensis*)
 RN 512-06-1 HCAPLUS
 CN Spirost-5-en-3-ol, (3 β ,25S)- (CA INDEX NAME)

Absolute stereochemistry.




IT 6869-59-6, 25L-Spirost-5-ene-3 β ,12 β -diol
 (as chiapagenin structure)
 RN 6869-59-6 HCAPLUS
 CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 6877-71-0, Diosgenin, 12 β -hydroxy-
 (as isochiapagenin structure)
 RN 6877-71-0 HCAPLUS
 CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 6869-59-6P, Chiapagenin
 RL: PREP (Preparation)
 (derivs., and separation from *Dioscorea chiapasensis*)
 RN 6869-59-6 HCAPLUS
 CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 512-04-9, Diosgenin 121193-55-3, Correllogenin, acetate
 (from *Dioscorea chiapasensis*)
 RN 512-04-9 HCAPLUS
 CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.




RN 121193-55-3 HCAPLUS
 CN 21 α ,22 α ,25L-Spirost-5-en-12-one, 3 β -hydroxy-, acetate
 (6CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 6877-71-0, Yamogenin, 12 β -hydroxy-
 (identity with chiapagenin)
 RN 6877-71-0 HCAPLUS
 CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 5996-01-0P, Botogenin, acetate 90457-38-8P,
 5 α ,25L-Spirostan-3 β ,12 β -diol 122677-90-1P,
 25L-Spirost-4-en-3-one, 12 β -hydroxy-
 RL: PREF (Preparation)
 (preparation of)
 RN 5996-01-0 HCAPLUS
 CN Spirost-5-en-12-one, 3-(acetyloxy)-, (3 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 90457-38-8 HCAPLUS
 CN Spirostan-3,12-diol, (3 β ,5 α ,12 β ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 122677-90-1 HCAPLUS

CN 25L-Spirost-4-en-3-one, 12 β -hydroxy- (6CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 59203-51-9P, Isochiapagenin


RL: PREP (Preparation)

(separation from *Dioscorea chiapasensis*, and its structure)

RN 59203-51-9 HCAPLUS

CN Spirost-5-ene-3,12-diol, (3 β ,12 β ,20 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 892496-15-0, Chiapagenin, dihydro-
(structure of)

RN 892496-15-0 HCAPLUS

CN INDEX NAME NOT YET ASSIGNED

Absolute stereochemistry.



L97 ANSWER 6 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1961:43404 HCAPLUS Full-text

DN 55:43404

OREF 55:8460a-i,8461a-i,8462a-i,8463a-i,8464a-b

TI The synthesis of the steroidal sapogenins

AU Mazur, Yehuda; Danieli, Naftali; Sondheimer, Franz

CS Weizmann Inst. Sci., Rehovoth, Israel

SO Journal of the American Chemical Society (1960), 82, 5889-908

CODEN: JACSAT; ISSN: 0002-7863

DT Journal

LA Unavailable

OS CASREACT 55:43404

GI For diagram(s), see printed CA Issue.

AB Isoandrosterone (I) was converted by an 18-stage process to a mixture of tigogenin (II) and neotigogenin (III). I treated with $\text{CH}_2\text{:CMeOAc}$ and H_2SO_4 , the resulting 16-androstene-3 β ,17-diol diacetate, 90%, m. 170-2°, treated with excess BzO_2H , and the crude epoxide treated 0.5 hr. with HClO_4 in AcOH at room temperature gave 60% 3 β ,16 α -diacetoxystrostan-17-one (IV), m. 183-5°, $[\alpha]_D^{25}$ 56° (all rotations in CHCl_3 unless noted otherwise). Activated granulated Zn (15 g.), 10 g. IV, and 100 cc. dry C_6H_6 heated to remove about 20 cc. C_6H_6 , cooled, treated with 40 g. $\text{MeCHBrCO}_2\text{Et}$ (V), the mixture heated a short time, refluxed 0.5 hr., decanted, the solution worked up, the residual yellow oil refluxed 3 hrs. in 200 cc. MeOH with 12 g. KOH in 10 cc. H_2O , diluted with H_2O , washed with Et_2O , acidified with dilute HCl , extracted with EtOAc , and the extract evaporated gave 3.5 g. 3 β ,16 α ,17 β -trihydroxy-17- isobisnor-5 α -cholanic acid (VI), m. 242-3° (MeOH), $[\alpha]_D^{25}$ -10° (dioxane). VI in CH_2Cl_2 treated 16 hrs. at 0° with $\text{CH}_2\text{N}_2\text{-Et}_2\text{O}$ gave Me ester (VII) of VI, m. 227-8° ($\text{Me}_2\text{CO-hexane}$), $[\alpha]_D^{25}$ -8° (dioxane). VI treated 16 hrs. at room temperature with $\text{Ac}_2\text{O-C}_5\text{H}_5\text{N}$ and heated 0.5 hr. at 90° with a little H_2O yielded 3,16-diacetate (VIII) of VI, m. 185-6° (MeOH), $[\alpha]_D^{25}$ -46° (dioxane). Acetylation of VII and treatment of VIII with $\text{CH}_2\text{N}_2\text{Et}_2\text{O}$ gave the Me ester of VIII, m. 172-3° (MeOH), $[\alpha]_D^{25}$ -37°. IV condensed in the usual manner with V and the product chromatographed on 500 g. Al_2O_3 yielded 0.72 g. Et 3 β ,16 α -diacetoxystrostan-17 β -hydroxy-17-isonor-5 α -cholanate (IX), m. 177-8° ($\text{Et}_2\text{O-hexane}$), $[\alpha]_D^{25}$ -38°, 0.35 g. 3 β -acetoxystrostan-16 α ,17 β -dihydroxy-17-isonor-5 α -cholanic 22 \rightarrow 16-lactone (X), m. 220-2° (MeOH), $[\alpha]_D^{25}$ -55°, 3.92 g. 16 α -OH analog (XI) of IX, m. 167-8° ($\text{Me}_2\text{CO-hexane}$), $[\alpha]_D^{25}$ -14°, 0.42 g. 3 β -OH analog (XII) of IX, m. 154-5° ($\text{Me}_2\text{CO-hexane}$), and 2.45 g. 3 β ,16 α -di-OH analog (XIII) of IX, m. 201-3° (Me_2CO), $[\alpha]_D^{25}$ -5°. Acetylation of XI, XII, and XIII gave IX, m. 176-8°. VIII (200 mg.) in 50 cc. dry Et_2O and 2 cc. SOCl_2 kept 2 hrs., evaporated in vacuo, and the residue refluxed 30 min. with 10 cc. absolute EtOH yielded 115 mg. IX, m. 176-7° ($\text{Et}_2\text{O-hexane}$). IX (200 mg.) in 2:1 pentane- C_6H_6 chromatographed on 10 g.

Al₂O₃ yielded 55 mg. unchanged IX and 125 mg. XI, m. 165-7°. XII gave similarly 60% XIII, m. 200-2°, but X, XI, and XIII were recovered unchanged under the same conditions. VI (100 mg.) in 25 cc. Ac₂O refluxed 2 hrs., evaporated in vacuo, the residue boiled 15 min. with H₂O, and the product isolated with Et₂O gave 62 mg. X, m. 218-20° (MeOH), [α]_D -56°. VI (100 mg.) in 20 cc. glacial AcOH treated 1 hr. at room temperature with dry HCl, the mixture poured into iced H₂O, and the product isolated with Et₂O gave 46 mg. X. VIII (100 mg.) and 200 mg. KHSO₄ heated 10 min. at 170°/about 1 mm. yielded 62 mg. X. X (200 mg.) and 1.5 g. KOH in 50 cc. 95% MeOH refluxed 2 hrs., treated with H₂O and Et₂O, the aqueous phase acidified with dilute HCl, extracted with EtOAc, the extract washed (aqueous Na₂CO₃ and H₂O), evaporated, and the residual 3-OH analog (31 mg.) of X acetylated gave X; the aqueous alkaline washing acidified with HCl and the product isolated with EtOAc gave 137 mg. VI, m. 240-2° (Me₂CO). IV (20 g.) condensed in the usual manner with 80 g. V, the crude product treated 16 hrs. at room temperature with 50 cc. Ac₂O and 50 cc. C₅H₅N, the product isolated with Et₂O, and chromatographed on 750 g. Al₂O₃ yielded 3.8 g. IX, m. 177-8° (Et₂O-hexane), and 11.3 g. XI, which acetylated gave 12.1 g. IX. 3β,17β-Diacetoxyandrostan-16-one (2.5 g.), m. 180-1°, [α]_D -118°, treated with 4 g. activated Zn and 10 g. V in 25 cc. C₆H₆ and the product chromatographed on 100 g. Al₂O₃ gave 0.91 g. 3-acetate of 3β,16,17β-trihydroxy-16- [1-(carbethoxy)ethyl] androstane isomer A (XIV), m. 188-9° [α]_D 5°, (diacetate m. 174-5°, [α]_D 2°), and 0.42 g. 3-acetate of isomer B, m. 123-4°, [α]_D -3° (diacetate m. 125-6°, [α]_D -14°). VIII (200 mg.), 5 cc. SOCl₂, and 5 cc. dry C₆H₆ refluxed 2 hrs., evaporated, the residue dissolved in 5 cc. C₆H₆, the solution added under N to Me₂Cd from 620 mg. MeI, 100 mg. Mg, 10 cc. Et₂O, and 100 mg. CdCl₂, the mixture stirred 2 hrs. under N at room temperature, kept 16 hrs., worked up, and the crude product chromatographed on 12 g. Al₂O₃ yielded 82 mg. 3β, 16α-diacetoxy-17β-hydroxy-17-isobisnor-5α-cholan-22-one, m. 190-1°, [α]_D -62°, unchanged upon heating with dioxane and H₂SO₄, but gave oily products when heated with POCl₃ and C₅H₅N or with Ac₂O. IX (1 g.) and 2 g. KHSO₄ heated 15 min. at 170-5°/about 25 mm., cooled, treated with Et₂O, the Et₂O phases from 3 runs worked up, and the crude product chromatographed on 150 g. Al₂O₃ gave 0.33 g. oily material, 1.15 g. Et 3β,16α- diacetoxybisnor-5α-chol-17(20)-enate (XV), m. 140-1° (Et₂O-hexane), [α]_D -73°, and 0.90 g. 3β-acetoxy- 16α-hydroxybisnor-5α-chol-17(20)-enic 22 → 16-lactone (XVI), m. 239-40° (Me₂CO-hexane), [α]_D -165°. A similar run with a longer reaction time gave more XVI at the expense of XV. IX (100 mg.) and 300 mg. CuSO₄ heated 0.5 hr. at 180°/25 mm. and the product chromatographed yielded 82 mg. oil and 7 mg. XV. IX (100 mg.) in 5 cc. C₅H₅N heated 0.5 hr. on the water bath with 3 cc. POCl₃ in 5 cc. C₅H₅N gave 80 mg. oil. IX (200 mg.) in 30 cc. Ac₂O refluxed 2 hrs., evaporated, and the residue chromatographed on 5 g. Al₂O₃ gave 175 mg. oil. XV (100 mg.) and 200 mg. KHSO₄ heated 0.5 hr. at 175°/25 mm. and the crude product chromatographed on 5 g. Al₂O₃ yielded 36 mg. XV and 24 mg. XVI, m. 236-9°. XV (120 mg.) and 450 mg. KOH in 15 cc. 95% MeOH refluxed 2 hrs., treated with H₂O and Et₂O, the aqueous alkaline solution acidified with dilute HCl, and the product isolated with Et₂O yielded 24 mg. XVI, m. 236-9° (Me₂CO-hexane). XI (250 mg.) and 500 mg. KHSO₄ heated 0.5 hr. at 170-80°/20 mm. and the product chromatographed on 12 g. Al₂O₃ yielded 103 mg. lactone, C₂₄H₃₄O₄ (XVII), needles, m. 165-6° (Et₂O-hexane), [α]_D -49°, and 78 mg. hydroxylactone, C₂₄H₃₆O₅, m. 250-2° (Me₂CO-hexane), [α]_D -65°. XVII (80 mg.), 300 mg. KOH, and 10 cc. 90% MeOH refluxed 2 hrs., diluted with Et₂O and H₂O, the aqueous alkaline layer acidified with dilute HCl, and the product isolated with EtOAc gave 62 mg. lactone, m. 175-7°, which reacetylated yielded XVII. XV (2 g.) in 60 cc. glacial AcOH hydrogenated 4 hrs. at 24°/764 mm. gave 1.71 g. Et 3β,16α-diacetoxy-20-isobisnor-5α-cholanate (XVIII), m. 129-30° (hexane), [α]_D -51°; 2nd polymorphic modification m. 159-60°. XVIII (1.5 g.) and 15 g. KOH in 150 cc. 85% EtOH refluxed 8 hrs., treated with H₂O and Et₂O, the aqueous layer

washed with Et₂O, acidified with dilute HCl, and the product isolated with EtOAc gave 1.13 g. 3 β ,16 α -dihydroxy-20-isobisnor-5 α -cholanic acid (XIX), which in 100 cc. absolute MeOH treated 16 hrs. at 0° with excess CH₂N₂-Et₂O gave 0.97 g. Me ester (XX), m. 181-2° (Me₂CO-hexane), [α]_D -7°; diacetate of XX m. 169-70°, [α]_D -50°. XVIII (200 mg.), 750 mg. KOH, and 25 cc. 95% MeOH refluxed 2 hrs. gave 76 mg. XX; the aqueous alkaline layer from the processing procedure acidified and extracted with EtOAc gave 68 mg. XIX, m. 180-2°. XVIII (0.5 g.) in 125 cc. tetrahydrofuran reduced with 1.25 g. LiAlH₄ in 60 cc. Et₂O gave 0.32 g. 20-isobisnor-5 α -cholane-3 β ,16 α ,22-triol (XXI), m. 269-70° (MeOH), [α]_D -23° (C₅H₅N); triacetate m. 151-2° (Et₂O-hexane), [α]_D -57°. XX (50 mg.) in 10 cc. tetrahydrofuran with 125 mg. LiAlH₄ in 10 cc. Et₂O yielded 35 mg. XXI. XX (1.1 g.) in 100 cc. glacial AcOH treated at 10° during 10 min. with 1 g. CrO₃ in 10 cc. 90% AcOH, kept 1 hr. at 10° and 1 hr. at room temperature, and worked up yielded 0.74 g. Me 3,16-dioxo-20-isobisnor-5 α -cholanate (XXII), m. 143-5° (EtAc-hexane), [α]_D -110°. XXI (300 mg.) in 25 cc. AcOH with 250 mg. CrO₃ in 2.5 cc. 90% AcOH gave 175 mg. acidic material, which in 20 cc. CH₂Cl₂ treated with CH₂N₂-Et₂O during 16 hrs. at 0° gave 140 mg. XXII. XXII (100 mg.) in 20 cc. MeOH treated 16 hrs. at room temperature with 500 mg. NaBH₄ in 4 cc. MeOH, the mixture worked up, the product (90 mg.) treated 16 hrs. at room temperature with 4 cc. C₅H₅N and 2 cc. Ac₂O, separated into 17 mg. acidic and 76 mg. neutral material, and the latter chromatographed on 5 g. Al₂O₃ yielded 48 mg. 3 β -acetoxy-16 β -hydroxy-20-isobisnor-5 α -cholanic 22 \rightarrow 16-lactone (20-isotigonein lactone acetate) (XXIII), m. 226-8° (hexane), [α]_D -36°. Crude XIX (100 mg.), 1 cc. concentrated HCl, and 1 cc. H₂O in 20 cc. glacial AcOH refluxed 2 hrs., poured into ice, the mixture extracted with EtOAc, the extract worked up, and chromatographed on 5 g. Al₂O₃ yielded 58 mg. XXIII, m. 224-6°, [α]_D -35°. XXIII (60 mg.) and 750 mg. KOH in 25 cc. 90% MeOH refluxed 2 hrs., washed with EtOAc, acidified, the product isolated with EtOAc, and acetylated gave 56 mg. 20-normal isomer (tigogenin lactone acetate) (XXIV) of XXIII, m. 219-21° (CH₂Cl₂-hexane), [α]_D -49°. XXIII was also rearranged to XXIV with NaOMe in C₆H₆ at 78° in a sealed tube. XVI (500 mg.) in 20 cc. EtOAc hydrogenated 1.5 hrs. at 27°/754 mm. over 50 mg. prerduced PtO₂ gave 445 mg. 3 β -acetoxy-16 α -hydroxy-17-isobisnor-5 α -cholanic 22 \rightarrow 16-lactone (XXV), m. 199-201° (CH₂Cl₂-hexane), [α]_D 21°. XXV (200 mg.) in 50 cc. tetrahydrofuran and 500 mg. LiAlH₄ in 20 cc. Et₂O gave 145 mg. 17-isobisnor-5 α -cholane-3 β ,16 α ,22-triol (XXVI), m. 245-7° (MeOH), [α]_D -45° (C₅H₅N); triacetate m. 100-1° (Et₂O-hexane), [α]_D -45°. XXVI (200 mg.) in 25 cc. AcOH oxidized with 200 mg. CrO₃ in 2 cc. 90% AcOH, worked up in the usual manner, the acidic fraction (140 mg.) in 20 cc. CH₂Cl₂ treated 16 hrs. at 0° with CH₂N₂-Et₂O, and the product chromatographed on 10 g. Al₂O₃ yielded 115 mg. Me 3,16-dioxo-17-isobisnor-5 α -cholanate (XXVII), m. 131-3° (CH₂Cl₂-hexane), [α]_D -67°. XXV (200 mg.) in 50 cc. 90% MeOH refluxed 2 hrs. and worked up with H₂O and Et₂O gave 175 mg. 3 β -acetoxy-16 α -hydroxy-17-iso-20-isobisnor-5 α -cholanic 22 \rightarrow 16-lactone (XXVIII), m. 190-1° (Me₂CO-hexane), [α]_D 13°. XVIII (200 mg.) in 50 cc. tetrahydrofuran and 500 mg. LiAlH₄ in 20 cc. Et₂O yielded 135 mg. 17-iso-20-isobisnor-5 α -cholane-3 β ,16 α ,22-triol (XXIX), m. 189-91° (MeOH-EtOAc), [α]_D -19° (C₅H₅N); triacetate m. 109-10° (Et₂O-hexane), [α]_D -71°. XXIX (100 mg.) in 10 cc. AcOH oxidized with 100 mg. CrO₃ in 1 cc. 90% AcOH, worked up, and the acidic product (48 mg.) chromatographed on 2 g. Al₂O₃ yielded 38 mg. Me 3,16-dioxo-17-iso-20-isobisnor-5 α -cholanate (XXX), m. 121-3°, [α]_D -121°. Natural XXIV reduced with LiAlH₄ gave about 80% bisnor-5 α -cholane-3 β ,16 β ,22-triol (XXXI), m. 246-9°, [α]_D 15° (C₅H₅N); triacetate m. 117-18° (Et₂O-hexane), [α]_D 52°. XXII (700 mg.) and 9 g. KOH in 300 cc. 93% aqueous MeOH refluxed 2 hrs., worked up in the usual manner, and the acidic material (620 mg.) in 100 cc. CH₂Cl₂ treated 16 hrs. at 0° with CH₂N₂-Et₂O yielded 640 mg. Me 3,16-dioxobisnor-5 α -cholanate (XXXII), m. 219-22° (EtCOMe),


[α]_D -108°. XXVII (100 mg.) treated with base and reacylated yielded 88 mg. XXXII, m. 214-19°. XXX (20 mg.) gave similarly 14 mg. XXXII, m. 218-20°. XXXI oxidized with CrO₃ in AcOH and the acidic product esterified with CH₂N₂-Et₂O gave about 40% XXXII, m. 220-2°, [α]_D -108°. XXXII (1 g.), 1.5 g. p-MeC₆H₄SO₃H.H₂O, 12 cc. (CH₂OH)₂, and 1.5 l. C₆H₆ refluxed 5 hrs. with the removal of 200 cc. C₆H₆, treated again with 1.5 g. p-Me₂CH₄SO₃H, refluxed 6 hrs. with the removal of 300 cc. C₆H₆, treated 10 min. with 3 g. NaOH in 50 cc. 95% MeOH, worked up, and the crude product chromatographed on 60 g. Al₂O₃ yielded 0.72 g. Me 3,3:16,16-bis(ethylenedioxy)bisanor-5 α -cholanate (XXXIII), m. 196-8° (Me₂CO-hexane), [α]_D -24°. XXXIII (1.13 g.) in 300 cc. tetrahydrofuran treated dropwise during 15 min. with 1.5 g. LiAlH₄ in 75 cc. Et₂O under N, the mixture stirred 1 hr., kept 16 hrs., and worked up gave 0.76 g. 3,3:16,16-bis(ethylenedioxy)bisanor-5 α -cholan-22-ol (XXXIV), m. 235-7° (Me₂CO), [α]_D -18°. XXXIV (40 mg.) in 10 cc. 90% AcOH heated 1 hr. at 90°, diluted with H₂O, the product dissolved in 10 cc. Et₂O, and reduced with 50 mg. LiAlH₄ in 2 cc. Et₂O gave 26 mg. XXXI, m. 247-50° (MeOH), [α]_D 14° (C₅H₅N). XXXIV (500 mg.) in 15 cc. C₅H₅N added dropwise to 350 mg. CrO₃ in 15 cc. dry C₅H₅N, the mixture kept 4 hrs. at 37°, worked up, the crude product isolated with EtOAc, and chromatographed on 30 g. Al₂O₃ yielded 415 mg. 3,3:16,16-bis(ethylenedioxy)bisanor-5 α -cholan-22-al (XXXV), m. 183-4° (Et₂O-pentane), [α]_D -21°. XXXV (50 mg.) in 10 cc. tetrahydrofuran reduced with 50 mg. LiAlH₄ in 2 cc. Et₂O at room temperature gave 41 mg. XXXIV, m. 235-7°. XXXV (80 mg.) in 7.5 cc. C₆H₆ added dropwise with stirring to iso-AmMgBr from 18 mg. Mg and 122 mg. iso-AmBr in 7.5 cc. Et₂O under N, the mixture refluxed 4 hrs., and worked up gave 67 mg. 3,3:16,16-bis(ethylenedioxy)cholestan-22-ol (XXXVI), m. 195-6° (Me₂CO-hexane), [α]_D -20°. XXXVI (40 mg.) in 2 cc. C₅H₅N treated with 40 mg. CrO₃ in 2 cc. C₅H₅N, the mixture kept 48 hrs. at 37°, worked up, and the product chromatographed on Al₂O₃ yielded 27 mg. 22-one analog (XXXVII) of XXXVI, m. 134-5° (Et₂O-pentane), [α]_D -16°. XXXVII (20 mg.) in 5 cc. 80% AcOH heated 1 hr. at 90°, the product isolated with Et₂O, and chromatographed on 3 g. Al₂O₃ gave 11 mg. cholestane-3,16,22-trione (XXXVIII), plates, m. 176-7°. XXXVIII (5 mg.) refluxed 1 hr. with 2 cc. 5% KOH-MeOH under N and the product isolated with Et₂O yielded XXXIX, oil. EtOCH:CHCO₂Et (108 g.) and 115 g. MeCH:CHCH₂OH containing 1 g. Na heated to 220° until the gas evolution ceased, the residue dissolved in 500 cc. MeOH, refluxed 2 hrs. with 50 g. NaOH in 100 cc. H₂O, diluted with H₂O, and worked up with Et₂O gave 25.8 g. CH₂:CHCHMeCH₂CO₂H (XL), b₂₅ 78-80°, n_{20D} 1.4366. XL (20 g.) in 200 cc. Et₂O treated dropwise with stirring with 8 g. LiAlH₄ in 400 cc. Et₂O, the mixture stirred 2 hrs., and worked up gave 13.8 g. CH₂:CHCHMeCH₂CH₂OH (XLI), b₂₅ 63-4°, n_{20D} 1.4369. XLI (12 g.) and 2.25 cc. C₅H₅ treated dropwise during 0.5 hr. with 4.5 cc. PBr₃, the mixture stirred 0.5 hr., and worked up gave 8.2 g. CH₂:CHCHMeCH₂CH₂Br (XLII), b₇₆₄ 138-40°, n_{20D} 1.4680. XXXV (300 mg.) in 10 cc. C₆H₆ added dropwise at room temperature with stirring to the Grignard reagent from 68 mg. Mg and 650 mg. XLII in 10 cc. Et₂O under N, the mixture refluxed 2 hrs., worked up, and the product chromatographed on 18 g. Al₂O₃ yielded 265 mg. mixed C-25 isomeric 3,3:16,16-bis(ethylenedioxy)-26-methylenecholestan-22-ol (XLIII), m. 143-57°, [α]_D -18°. CrO₃ (400 mg.) added slowly with cooling to 20 cc. C₅H₅N, the mixture kept 48 hrs. at 37° with 400 mg. XLIII, decomposed with MeOH, and the product isolated with EtOAc gave 340 mg. 22-one analog (XLIV) of XLIII, m. 145-8° (Et₂O), [α]_D -16°. XLIV (30 mg.) in 15 cc. 90% AcOH heated 1 hr. at 90°, the product isolated with Et₂O, and chromatographed on 4 g. Al₂O₃ gave 18 mg. 26-methylenecholestane-3,16,22-trione, m. 161-5° (Et₂O-pentane). XLIV (300 mg.) in 30 cc. EtOAc containing 3 drops of C₅H₅N ozonized at -18°, treated with 6 g. Raney Ni, refluxed 10 min., filtered, evaporated, the residue heated 0.5 hrs. at 90° with 30 cc. 80% AcOH, and worked up gave 195 mg. crude noncryst. aldehydes; the crude product dissolved in 90 cc. dry tetrahydrofuran, the solution reduced with 300 mg. NaBH₄ in 90 cc. absolute iso-PrOH during 72 hrs. at room temperature, the

product isolated with EtOAc, heated 5 min. on the water bath with 30 cc. MeOH and 0.3 cc. 10% aqueous HCl, and chromatographed on 9 g. Al₂O₃ yielded 63 mg. mixture of II and III, needles, m. 178-82° (sublimed at 160°/0.01 mm.), [α]_D -71°. II-III mixture (15 mg.) in 25 cc. EtOH and 6 cc. concentrated HCl refluxed 48 hrs. under N, diluted with 3 cc. concentrated HCl, refluxed 72 hrs., the product isolated with EtOAc, and chromatographed on Al₂O₃ gave 9 mg. II, m. 202-4° (Me₂CO-hexane), [α]_D -68°. II-III mixture (45 mg.) acetylated and the mixed acetates recrystd. from a relatively dilute EtOH solution gave 10 mg. acetate of III, octahedra, m. 175-8°, which refluxed 1 hr. with 20 cc. 3% KOH in 90% MeOH and worked up yielded 7 mg. III, m. 201-3° (Me₂CO-hexane), [α]_D -76°. 5α,25D-spirostan-3-one (XLV) (1 g.) in 150 cc. glacial AcOH treated with 1 cc. HBr-AcOH and then during 3 min. with stirring at room temperature with 1.3 g. Br in 20 cc. AcOH, the mixture kept 10 min., and worked up gave 0.88 g. 2α,4α,23-tribromo- 5α,25D-spirostan-3-one (XLVI), m. 196-8° (decomposition) (CH₂Cl₂-EtOAc).

Br (200 mg.) added to 15 cc. Me₂CO, the mixture treated with 1 g. Na₂CO₃, shaken 20 min., filtered, added to 5 g. NaI in 100 cc. Me₂CO, refluxed 0.5 hr., treated with 850 mg. XLVI, refluxed 12 hrs., worked up, the product refluxed 3 hrs. with 10 g. Zn dust in 100 cc. AcOH, worked up, and chromatographed on 40 g. Al₂O₃ yielded 280 mg. XLV and 165 mg. 25D-spirost-4-en-3-one (XLVII), m. 185-7° (CHCl₃-Et₂O), [α]_D -7°. XLVII (150 mg.) in 50 cc. CH₂:CMeCO₂Ac refluxed 3 hrs. and worked up gave 115 mg. 25D-spirosta-3,5-dien-3-ol acetate (XLVIII), m. 181-2° (Et₂O-MeOH), [α]_D -113°. XLVIII (100 mg.) in 200 cc. EtOH added dropwise during 2 hrs. with stirring to 1 g. NaBH₄ in 50 cc. 70% EtOH at 5°, the mixture kept 1 hr. at 5°, worked up, the product isolated with EtOAc, refluxed 1 hr. in 50 cc. EtOH with 3 drops concentrated HCl, again isolated with EtOAc, and chromatographed on 6 g. Al₂O₃ yielded 66 mg. diosgenin, m. 205-7°, (MeOH) [α]_D -119°. 5α,25D-Spirost-9(11)-en-3β-ol acetate (XLIX) (75 mg.), m. 197-8°, in 15 cc. AcOH treated 48 hrs. at 37° with 75 mg. CrO₃ in 5 cc. 85% AcOH, worked up, and the product chromatographed on 6 g. Al₂O₃ yielded 22 mg. XLIX and 18 mg. 3α-acetoxy-5α,25D-spirost-9(11)-en-12-one (L), m. 217-19° (MeOH), [α]_D -9°. L (50 mg.) in 20 cc. dry Et₂O added dropwise during 5 min. with stirring to 100 mg. Li in about 30 cc. liquid NH₃, the mixture stirred 5 min., worked up, the product refluxed 2 hrs. with 20 cc. 3% KOH-MeOH (containing 2 cc. H₂O), isolated with EtOAc, and chromatographed on 5 g. Al₂O₃ yielded 31 mg. hecogenin, m. 263-5° (Me₂CO), [α]_D 6°.

IT 77-60-1P, Tigogenin 470-01-9P, Neotigogenin
4948-43-0P, Neotigogenin, acetate 6870-79-7P,
25D-Spirost-4-en-3-one 6877-75-4P, 25D-Spirosta-3,5-dien-3-ol,
acetate 121250-54-2P, 5α,25D-Spirost-9(11)-en-12-one,
3α-hydroxy-, acetate
RL: PREP (Preparation)
(preparation of)
RN 77-60-1 HCAPLUS
CN Spirostan-3-ol, (3β,5α,25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 470-01-9 HCAPLUS

CN Spirostan-3-ol, (3 β ,5 α ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 4948-43-0 HCAPLUS

CN Spirostan-3-ol, acetate, (3 β ,5 α ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 6870-79-7 HCAPLUS

CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 6877-75-4 HCAPLUS

CN Spirosta-3,5-dien-3-ol, acetate, (25R)- (8CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 121250-54-2 HCAPLUS

CN 5 α ,25D-Spirost-9(11)-en-12-one, 3 α -hydroxy-, acetate (6CI)
(CA INDEX NAME)

Absolute stereochemistry.




IT 512-04-9P, Diosgenin

RL: PREP (Preparation)
(synthesis of)

RN 512-04-9 HCAPLUS

CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 7 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1960:103641 HCAPLUS Full-text

DN 54:103641

OREF 54:19759e-i,19760a-i,19761a-i,19762a-b

TI Long-range effects in alicyclic systems. III. Relative rates of condensation of some steroid and triterpenoid ketones with benzaldehyde

AU Barton, D. H. R.; McCapra, F.; May, P. J.; Thudium, F.

CS Univ. Glasgow, UK

SO Journal of the Chemical Society (1960) 1297-1311

CODEN: JCSOA9; ISSN: 0368-1769

DT Journal

LA Unavailable

AB cf. C.4 51, 13817b. The rates of alkali catalyzed condensation of a series of steroidal 3-ones with BzH (I) to give the corresponding 2-benzylidene derivs. were determined. As in the earlier work with triterpenoid ketones, long-range effects produced by unsatd. substituents (especially the ethylenic linkage) and by other groups could be easily detected. There existed a quant. relation between the rates for structurally analogous steroidal and triterpenoid ketones such that rates could be expressed in terms of the rate of a saturated reference ketone multiplied by a series of group rate factors (f) each of which was characteristic of the nature and position of the substituent group. The possible role of polar factors in influencing rates of condensation of carbonyl substituted ketones was admitted, but the major importance of the new effects of conformational transmission was considered to have been again demonstrated for ketones having remotely placed ethylenic substitution. A cursory investigation of derivs. of β -decalone was shown, that, wherever structurally appropriate, the same effects could be recognized and were of the same quant. magnitude as in corresponding steroid and triterpenoid ketones. A preliminary account of this work was given earlier (CA 53, 17875b). Lanostane-3,11-dione (1.16 g.), 0.22 ml. (CH₂OH)₂, 10 mg. p-MeC₆H₄SO₃H, and 50 ml. C₆H₆ refluxed 18 hrs., the solution poured into saturated aqueous Na₂CO₃, and the C₆H₆ layer separated gave 1.13 g. lanostane-3,11-dione 3-(ethylene ketal) (II), m. 140-1° (C₆H₆-MeOH), [α]_D 31° (c 1.37, all rotations refer to CHCl₃ unless otherwise specified). II (150 mg.) in 10 ml. refluxing PrOH treated during 1 hr. with 1 g. Na, 7 ml. PrOH added, the solvent removed, and the residue worked up as usual gave 131 mg. 11 α -hydroxylanostan-3-one ethylene ketal (III), prisms, m. 165-6° (MeOH), [α]_D 0° (c 1.88). Hydrolysis of 180 mg. III in 12 ml. AcOH and 3.5 ml. H₂O 10 min. on the steam bath gave 150 mg. 11 α -hydroxylanostan-3-one, prisms, m. 151-2° (aqueous MeOH), [α]_D -6° (c 1.66). II (200 mg.) in 50 ml. dry Et₂O refluxed 16 hrs. with 220 mg. LiAlH₄, the excess reducing agent destroyed with EtOAc, and the mixture worked up as usual gave 165 mg. 11 β -hydroxylanostan-3-one ethylene ketal (IV), m. 143-4° (MeOH), [α]_D 30° (C 2.00). IV (270 mg.) in 35 ml. AcOH and 5 ml. H₂O heated 10 min. gave 157 mg. 11 β -hydroxylanostan-3-one, plates, m. 188-9° (ligroine), [α]_D 28° (c 1.49). IV (279 mg.) treated 15 min. at room temperature with 12


drops aqueous 60% HClO₄ in 15 ml. AcOH gave 211 mg. lanost-9 (11)-en-3-one, m. 113-14° (CHCl₃-MeOH), [α]_D 65° (c 2.67). 3β-Hydroxy-α-amyr-12-en-11-one (1.5 g.) in 100 ml. C₆H₆ added dropwise to 200 ml. Et₂O containing MeMgI (from 8 ml. MeI), the Et₂O distilled, the C₆H₆ solution refluxed 55 hrs., excess of saturated aqueous NH₄Cl added, the C₆H₆ separated, the solvent evaporated, the residue left overnight at room temperature with 10 ml. C₅H₅N and 10 ml. Ac₂O and chromatographed on Al₂O₃ gave 752 mg. 11-methylene-α-amyr-12-enyl acetate (V), m. 229-32° (CHCl₃-MeOH), [α]_D 143° (c 1.95), λ 246 mμ, ε 19,700. Hydrolysis of V gave the alc. and oxidation with C₅H₅N-CrO₃ gave 11-methylene-α-amyr-12-en-3-one, m. 146-7° (aqueous MeOH), [α]_D 208° (c 1.2), λ 247 mμ, ε 19,700. β-Amyrane-3,12-dione (1.7 g.) in 160 ml. (CH₂OH)₂ containing 60 mg. p-MeC₆H₄SO₃H slowly distilled at 1.5 mm. during 2 hrs. and dilute aqueous KOH added gave 1.59 g. β-amyrane-3,12-dione 3-(ethylene ketal) (VI), m. 279-81° (C₆H₆-MeOH), [α]_D -45° (c 1.64). VI (150 mg.) in 10 ml. refluxing PrOH treated during 1 hr. with 1 g. Na gave 121 mg. 12β-hydroxy-β-amyr-3-one ethylene ketal (VII), plates, m. 272-4° (C₆H₆-MeOH). Treatment of VII (167 mg.) with 5 ml. 90% AcOH 5 min. at 100° gave 87 mg. 12β-hydroxy-β-amyr-3-one, m. 210-13° (aqueous MeOH), [α]_D 39° (c 1.54). VII (474 mg.) reduced with 900 mg. LiAlH₄ in 150 ml. refluxing Et₂O gave on chromatography 254 mg. 12α-hydroxy-β-amyr-3-one ethylene ketal (VIII), m. 261-3° (C₆H₆-MeOH), [α]_D 16° (c 2.13). Elution with C₆H₆ gave 123 mg. VII. VIII treated with aqueous AcOH gave 12α-hydroxy-β-amyr-3-one, m. 252-5° (CHCl₃-MeOH), [α]_D 81° (c 1.11). Wolff-Kishner reduction of 1.9 g. 12-oxo-β-amyranyl acetate gave after reacetylation 1.24 g. β-amyranyl acetate. Alkaline hydrolysis and oxidation with C₅H₅N-CrO₃ gave β-amyr-3-one, m. 200-1° (CHCl₃-MeOH), [α]_D 41° (c 0.97). 7-Oxocholestanyl acetate was converted into 80% 7-methylenecholestanyl acetate, leaflets, m. 72-3° (alc.), [α]_D -43° (c 0.95), ν 890 and 1650 cm.⁻¹ Refluxing 0.5 hr. with MeOH-KOH gave 7-methylenecholestanol (IX), m. 115° (ligroine), [α]_D -31° (c 1.14). IX (560 mg.) in 60 ml. Me₂CO treated with a standard solution of CrO₃ in concentrated H₂SO₄ under N with shaking 3 min., gave after extraction 530 mg. 7-methylenecholestanone, plates, m. 104-6° (aqueous alc.), [α]_D -11° (c 0.94), ν 1700, 890, and 1650 cm.⁻¹ Hecogenin acetate (1.5 g.) in 20 ml. C₆H₆ treated 1 hr. at room temperature with MeMgI in Et₂O, and warmed 45 min. at 40° gave 910 mg. 12β-hydroxy-12α-methyltigogenin (X), m. 197-9° (Et₂O), [α]_D -37° (c 1.00). X (720 mg.) in 70 ml. Me₂CO treated 7 min. at 0° with standard oxidation mixture gave 600 mg. 12β-hydroxy-12α-methyltigogenone (XI), plates, m. 228-43°, prisms, m. 241-3° (aqueous alc.), [α]_D -21° (c 1.06). XI (245 mg.) in 12 ml. C₅H₅N treated 16 hrs. at room temperature with POCl₃ and then 2.5 hrs. at 55° and the product chromatographed on Al₂O₃ gave 12-methylenetigogenone (XII), m. 219-21° (ligroine), [α]_D -5° (c 0.88), ν 890 and 1650 cm.⁻¹ XII (49 g.) in 50 ml. CH₂Cl₂ ozonized 1 hr. at -80° gave 12 mg. hecogenone. 11-Dehydrotigogenin oxidized with CrO₃ to 11-dehydrotigogenone, irregular plates, m. 169-74° (MeOH), [α]_D -20° (c 1.13). 9 (11)-Dehydrotigogenin oxidized to 9 (11)-dehydrotigogenone, m. 195-6.5° (MeOH), [α]_D -45° (c 0.98). Ergosta-7,14,22-trienol (250 mg.) in 38 ml. C₆H₆, 10 ml. Me₂CO, and 2 g. (iso-PrO)₃Al refluxed 8 hrs. gave 70 mg. ergosta-7,14,22-trien-3-one (XIII), m. 150-2° (MeOH), [α]_D -220° (c 0.96), λ 242 mμ, ε 9800. XIII was also prepared by CrO₃-Me₂COH₂SO₄ oxidation of ergosterol B₃, but the yield was only 20%. Ergost-22-ene-3,11-dione (720 mg.) in 200 ml. (CH₂OH)₂ containing 60 mg. p-MeC₆H₄SO₃H slowly distilled during 5 hrs. at 63°/1.5 mm., and the product treated with alc. KOH gave 640 mg. ergost-22-ene-3,11-dione 3-(ethylene ketal) (XIV), plates, m. 153-4° (MeOH), [α]_D 19° (c 2.02). XIV (200 mg.) in 15 ml. refluxing PrOH treated with 1.5 g. Na gave 180 mg. 11α-hydroxyergost-22-en-3-one ethylene ketal (XV), needles, m. 184-5° (MeOH), [α]_D -23° (c 1.5). XV (150 mg.)

hydrolyzed 10 min. with 80% aqueous AcOH gave 100 mg. 11 α -hydroxyergost-22-en-3-one, m. 142-4° (ligroine), [α]D -19° (c 1.59). XIV (150 mg.) reduced with excess LiAlH₄ in Et₂O gave 120 mg. 11 β -hydroxyergost-22-en-3-one ethylene ketal (XVI), m. 155-6° (aqueous MeOH), [α]D 0° (c 1.6). Hydrolysis of XVI with aqueous AcOH gave 60 mg. 11 β -hydroxyergost-22-en-3-one, m. 170-2° (ligroine), [α]D 12° (c 2.0). 3 β -Acetoxyergostane-7,11-dione hydrolyzed as usual gave 3 β -hydroxyergostane-7,11-dione (XVII), m. 177-9° (MeOH), [α]D -6° (c 2.73). Oxidation of XVII with CrO₃AcOH and C₆H₆ gave ergostane-3,7,11-trione, m. 186-8°, [α]D 16.7° (c 1.92). Similar oxidation of 3 β -hydroxyergost-22-en-7,11-dione gave ergost-22-en-3,7,11-trione, plates, m. 194-5° (MeOH), [α]D (c 2.28). Ergosta-8,22-dienol oxidized with CrO₃ in C₅H₅N to ergosta-8,22-dienone, plates, m. 168-70° (MeOH), [α]D 47° (c 0.30). 17 β -Hydroxyandrostan-3-one hexahydrobenzoate (0.8 g.) in 10 ml. C₆H₆ and 10 ml. MeOH treated with 5 mg. p-MeC₆H₄SO₃H gave 0.5 g. 3,3-dimethoxyandrostan-17 β -yl hexahydrobenzoate (XVIII), m. 130-2°, [α]D 12° (c 2.03). XVIII reduced with LiAlH₄ gave 0.3 g. 3,3-dimethoxyandrostan-17 β -ol, m. 180-2° (aqueous MeOH), [α]D 14° (c 1.68). The ketal (0.3 g.) in 10 ml. MeOH and 2 ml. 4N H₂SO₄ left 1 hr. at room temperature, poured into H₂O, extracted with Et₂O, and processed as usual gave 0.2 g. 17 β -hydroxyandrostan-3-one, m. 178-9° (aqueous MeOH), [α]D 32° (c 1.91). This procedure gave a better yield than hydrolysis of the hexahydrobenzoate in the presence of the 3-one. 3 β -Hydroxy-11-oxobisnorallocholanolic acid (0.8 g.) in 50 ml. C₆H₆, oxidized with a slight excess of CrO₃ in aqueous AcOH gave 0.6 g. 3,11-dioxobisnorallocholanolic acid, plates, m. 258-61° (alc.), [α]D 52° (c 2.54). Stigmastanone (0.5 g.) in 50 ml. 0.1N alc. KOH treated 24 hrs. at room temperature in the dark with 0.5 g. I gave 310 mg. 2-benzylidenestigmastanone (XIX), m. 151-2° (MeOH-C₆H₆), [α]D -108° (c 1.46), λ 294 m μ , ϵ 16,200. Addition of H₂O to the mother liquor and extraction with Et₂O gave 80 mg. more material. XIX (563 mg.) in 100 ml. CHCl₃ at -60° ozonized during 20 min., 5 ml. H₂O added, the CHCl₃ evaporated, and the oil dissolved in 2% aqueous KOH, washed, and acidified gave 50% 2,3-secostigmastane-2,3-dioic acid, plates, m. 230-2° (C₆H₆), [α]D 33° (c 0.98). Ergost-22-ene-3,11-dione (0.3 g.) in 25 ml. 0.1N alc.-KOH treated 24 hrs. with 300 mg. I at room temperature in the dark gave 190 mg. 2-benzylideneergost-22-ene-3,11-dione, plates, m. 191-2° (alc.), [α]D -7° (c 1.52), λ 294 m μ , ϵ 17,000. 3,11-Dioxobisnorallocholanolic acid (0.2 g.) in 25 ml. 0.1N alc.-KOH treated as above with 0.3 g. I, poured into H₂O, acidified, and extracted gave 150 mg. 2-benzylidene-3,11-dioxobisnorallocholanolic acid, m. 268-70° (MeOH-C₆H₆), [α]D -24° (c 2.28), λ 294 m μ , ϵ 16,800. 7-Methylencholestanone (100 mg.) similarly yielded 65 mg. 2-benzylidene-7-methylencholestanone, m. 145-7° (MeOH-C₆H₆), [α]D -178° (c 1.16), λ 294 m μ , ϵ 17,700. Ergost-8(14)-enone similarly gave 2-benzylideneergost-8(14)-en-3-one, m. 162-3° (C₆H₆-MeOH), [α]D -18° (c 2.4), λ 294 m μ , ϵ 17,000. 17-Hydroxyandrostan-3-one (65 mg.) with I as above gave 50 mg. 2-benzylidene-17-hydroxyandrostan-3-one, m. 190-1° (MeOH), [α]D -140° (c 1.8), λ 294 m μ , ϵ 16,600. 17 β -Hydroxy-4,4-dimethylandrost-5-en-3-one (100 mg.) converted into 50 mg. 2-benzylidene-17 β -hydroxy-4,4-dimethylandrost-5-en-3-one, prisms, m. 159-61° (MeOH-C₆H₆), [α]D -148° (c 1.45), λ 294 m μ , ϵ 16,500. 4,4-Dimethylergosterone (0.3 g.) in 50 ml. tetrahydrofuran and 100 ml. EtNH₂ at 0° treated with Li, the solvent removed in vacuo, the residue oxidized at 0° with CrO₃ in Me₂CO, and the product chromatographed on Al₂O₃ gave 150 mg. product, m. 176-80°, [α]D -33° (c 1.36). Further elution gave 100 mg. 4,4-dimethylergosta-7,22-dien-3-one (XX), m. 143-5° (MeOH), [α]D -37° (c 1.36). XX (50 mg.) treated with I gave 25 mg. 2-benzylidene-4,4-dimethylergosta-7,22-dien-3-one (XXI), m. 133-5° (MeOH-C₆H₆), [α]D -111° (c 1.9), λ 289 m μ , ϵ 17,300. I condensed with ergosta-7,22-dien-3-one gave 800 mg. oil derivative, λ 294 m μ , ϵ 13,500. This taken up in 10 ml.

C₆H₆ refluxed 14 hrs. in a solution of 200 mg. K in 10 ml. tert-BuOH and 5 ml. MeI gave 250 mg. XXI. Cholestanone (100 mg.) in 20 ml. 0.1N MeOH-KOH treated at room temperature with 100 mg. I gave 80 mg. 2-(α -hydroxybenzyl)cholestanone, m. 188-90° (MeOH-C₆H₆), [α]_D -71° (c 1.12). On treatment with alc. KOH under the conditions of a kinetic run this afforded in 5 min. I, 90% cholestanone, and 10% benzylidenecholestanone. Treatment of 50 mg. of the ketol with 20 ml. N alc. HCl gave the benzylidene derivative, amorphous, λ 294 m μ , ϵ 16,000. trans- β -Decalone (0.7 g.) in 25 ml. 0.1N alc. KOH treated 30 hrs. at room temperature in the dark with 2.6 g. I, and working as in earlier examples and trituration with ligroine gave 405 mg. 3-benzylidene-trans- β -decalone (XXII), prisms, m. 92-3° (ligroine), λ 292 m μ , ϵ 17,400. XXII (351 mg.) in 100 ml. CHCl₃ was ozonized 0.5 hr. at -20° until the absorption at 292 m μ disappeared. The solution worked up as above gave 250 mg. trans-cyclohexylidene-1,2-diacetic acid, prisms, m. 164-5°. Benzylidene derivative of stigmastanone (21.2 mg.), 19.8 mg. benzylidene of ergost-8(14)-en-3-one, and 10.4 mg. XXII was treated with a 10 molar excess of I in 0.1N alc. KOH; in 20 hrs. there was no change in the intensity of the ultraviolet absorption and no increase in the 330 m μ region. 2 α -Methylcholestanone (19.8 mg.) treated with I in alc. KOH as above gave 12 mg. unchanged material. The following ketones were treated under the conditions of a kinetic run: 3-hydroxycholestan-7-one, hecogenin, 3 β -hydroxyergost-22-ene-7,11-dione, and 3 β -hydroxyergost-22-en-11-one. In each case there was no appearance of ultraviolet absorption and the ketone was recovered unchanged.


IT 2137-26-4P, Hecogenone 7361-26-4P, Tigogenone,
 9(11)-dehydro- 15401-31-7P, Tigogenin, 12 β -hydroxy-12-
 methyl- 16127-92-7P, Tigogenone, 11-dehydro-
 117917-08-5P, Tigogenone, 12 β -hydroxy-12-methyl-
 RL: PREP (Preparation)
 (preparation of)
 RN 2137-20-4 HCAPLUS
 CN Spirostan-3,12-dione, (5 α ,25R)- (CA INDEX NAME)

Absolute stereochemistry.



RN 7361-26-4 HCAPLUS
 CN Spirosta-4,9(11)-dien-3-one, (25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 15401-31-7 HCAPLUS

CN Spirostan-3,12-diol, 12-methyl-, (3 β ,5 α ,12 β ,25R)- (9CI)
(CA INDEX NAME)


Absolute stereochemistry.



RN 16127-92-7 HCAPLUS

CN Spirost-11-en-3-one, (5 α ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 117917-08-5 HCAPLUS

CN Tigogenone, 12 β -hydroxy-12-methyl- (6CI) (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 8 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1960:28947 HCAPLUS Full-text

DN 54:28947

OREF 54:5738i,5739a-i,5740a-e

TI Steroidal components of domestic plants. XIX. Structure of kogagenin, a sapogenin from *Dioscorea tokoro*

AU Takeda, Kenichi; Kubota, Tokuo; Shimaoka, Ariyoshi

CS Shionogi & Co., Ltd., Osaka

SO Tetrahedron (1959), 7, 62-9

CODEN: TETRAB; ISSN: 0040-4020

DT Journal

LA Unavailable


AB cf. C.A. 53, 16206f. Kogagenin (I), a steroidal sapogenin isolated from the epigeous part of *D. tokoro*, was the 1st example of a naturally occurring spirostan tetrol. I, C₂₇H₄₄O₆, [α]_D -27° (C₅H₅N), m. 318-22° (decomposition), λ 10.90, 11.10 μ , (600 mg.) refluxed 2 hrs. in 6 ml. Ac₂O and 3 ml. C₅H₅N and the product isolated with C₆H₆ gave I triacetate (II), m. 249-52°, [α]_D -26° (c 1.0, CHCl₃). I was a 25D-sapogenin and since II still showed an infrared absorption OH band, I was assumed to be a 25D-tetrahydroxyspirostan. I (500 mg.) refluxed 22 hrs. in 500 ml. Me₂CO and 200 ml. C₆H₆ with 500 mg. p-MeC₆H₄SO₃H and the neutralized (Na₂CO₃) solution concentrated in vacuo, extracted with C₆H₆ and the washed and dried extract evaporated, the crystalline residue chromatographed on 15 g. Al₂O₃, and eluted with 1:1 C₆H₆-CHCl₃ and CHCl₃ gave 316 mg. material, recrystd. (CHCl₃-MeOH) to give 260 mg. I acetone (III), m. 273-5°, [α]_D -23° (c 1.15, CHCl₃). Further elution with 1:1 CHCl₃-MeOH gave 230 mg. I. II (500 mg.) in 5 ml. C₅H₅N at 0° treated dropwise with 0.50 g. SOCl₂ in 2 ml. C₅H₅N and the mixture kept 1 hr. at 0°, diluted with ice H₂O and extracted with Et₂O, and the washed (dilute HCl, aqueous NaHCO₃, H₂O) and dried extract evaporated yielded 490 mg. oil, crystallized (MeOH) to give 395 mg. anhydrokogagenin triacetate (IV), m. 171-3°, [α]_D 33° (c 1.0, CHCl₃). IV (500 mg.) refluxed 1 hr. in 20 ml. 1.5% KOH-MeOH and the cooled mixture diluted with H₂O, filtered, and the precipitate recrystd. (MeOH) gave anhydrokogagenin (V), m. 240-3°, [α]_D -70° (c 1.0, CHCl₃), neg. Rosenheim test. II was not affected by CrO₃C₅H₅N oxidation and the ready dehydration to IV showed I to have a cis- α glycol group and a tertiary OH function. IV (100 mg.) in 7 ml. AcOH hydrogenated 30 min. with 100 mg. prerduced PtO₂ and the filtered solution evaporated gave 45 mg. authentic tokorogenin triacetate (VI), m. 253-5°, [α]_D -20° (c 1.0, CHCl₃), saponified with 3% KOH-MeOH to tokorogenin, m. 266-8° (MeOH). The mother liquors from VI concentrated and the residue chromatographed on 3 g. Al₂O₃, the column washed free from 5 mg. VI with 1:1 petr. ether-C₆H₆, and eluted with C₆H₆ and 19:1 C₆H₆Et₂O gave 35 mg. dihydrotokorogenin triacetate (VII), m. 167-9°, [α]_D 40° (c 1.0, CHCl₃), λ 2.83 μ , but no spiroketal bands at 10.19, 10.90, 11.10, 11.58 μ , identical with a specimen prepared by catalytic

reduction of authentic VI. Accordingly, I was established as a hydroxytokorogenin. IV (300 mg.) in 5 ml. C₅H₅N and 350 mg. OsO₄ in 10 ml. C₆H₆ kept 53 hrs. in the dark at room temperature and the mixture saturated with H₂S, filtered, and the filtrate evaporated in vacuo yielded 150 mg. diol triacetate (VIII), m. 252-4°, [α]_D -44° (c 1.0, CHCl₃). VIII (100 mg.) in 10 ml. AcOH kept overnight at room temperature with 0.7 g. Pb(OAc)₄ in 20 ml. AcOH and diluted with H₂O, extracted with Et₂O, and the washed and dried ext evaporated produced a gum, λ 5.85 μ (strong), giving a pos. triphenyltetrazolium test, showing the presence of a CHO group and limiting the position of the double bond in IV to Δ 5 (or Δ 4) or Δ 14. IV (450 mg.) in 2 ml. C₅-H₅N and 2 ml. Ac₂O refluxed 5.5 hrs. with 0.3 g. EtNH₂.HCl and the cooled mixture poured onto ice, extracted with Et₂O, the furostene taken up in 9 ml. AcOH and treated dropwise with 0.3 g. CrO₃ in 3 ml. 80% AcOH, the mixture stirred 2 hrs. at room temperature and diluted with H₂O, extracted with Et₂O and the washed and dried extract evaporated, the gummy solid saponified with 1% alc. KOH and the product reacetylated, purified by chromatography, and recrystd. (dilute alc.) yielded 95 mg. 5, 16-pregnadiene-1 β ,2 β ,3 α -triol-20-one triacetate (IX), m. 150-2°, [α]_D 168° (c 1.0, CHCl₃), λ 239 m μ (log ϵ 4.00, alc.), establishing the presence of a Δ 16-20-ketone group without addnl. conjugation. V (100 mg.) refluxed 6 hrs. in 30 ml. Me₂CO containing 10 mg. p-MeC₆H₄SO₃H and the neutralized (NaHCO₃) solution concentrated in vacuo, extracted with Et₂O, the product chromatographed, and eluted with 9:1 petr. ether-C₆H₆ and with 4:1-1:1 petr. ether-C₆H₆ gave the diene acetonide (X), m. 162-4° (MeOH), [α]_D -115° (c 1.0, CHCl₃), λ 236 m μ (log ϵ 4.31, alc.), and 93 mg. 25D-spirost-5-ene-1 β ,2 β ,3 α -triol 1,2-acetonide (XI), m. 208-10° (MeOH), [α]_D -61° (c 1.0, CHCl₃). XI (80 mg.) in 4 ml. C₅H₅N containing 0.5 g. POCl₃ heated 45 min. on a steam bath and the solution poured onto crushed ice, extracted with Et₂O, and the oily product crystallized (MeOH) yielded 15 mg. X. Attempts to obtain a Δ 4-3-oxo derivative of I by oxidation of XI with CrO₃C₅H₅N complex, with CrO₃-Me₂CO-H₂SO₄, or by Oppenauer oxidation were unsuccessful with almost quant. recovery of XI. III (120 mg.) in 3 ml. C₅H₅N added at 0° to the complex from 150 mg. CrO₃ and 1.5 ml. C₅H₅N and the mixture kept 16 hrs. at room temperature, extracted with Et₂O, and the product crystallized (MeOH) gave a crude ketone (XII), m. 189-91° (decomposition), λ 2.84, 5.76 μ , contaminated by 10% α,β -unsatd. ketone (XIII). XII (80 mg.) chromatographed in 1:1 petr. ether-C₆H₆ on SiO₂ gel and eluted with 9:1 C₆H₆-CHCl₃ yielded 54 mg. material, m. 193-8°, λ 246 m μ (log ϵ 4.10, alc.), recrystd. (MeOH) to give XIII, m. 197-200°, [α]_D -100° (c 1.0, CHCl₃), λ 246 m μ (log ϵ 4.15, alc.), λ 5.93, 6.16 μ , but no OH absorption. XII (20 mg.) in 5 ml. MeOH kept overnight with 0.5 ml. 10% aqueous KOH and diluted with H₂O, neutralized with dilute HCl, and extracted with Et₂O yielded 15 mg. 25D-spirosta-1,4-dien-2-ol-3-one (XIV), m. 224-7°, [α]_D -102° (c 0.36, CHCl₃), λ 2.96, 6.09, 6.17 μ (Nujol), λ 254 m μ (log ϵ 4.13, alc.), giving reddish purple color with alc. FeCl₃. XIV refluxed with alc. o-(H₂N)₂C₆H₄ produced an orange-yellow quinoxaline. I (98 mg.) in 4 ml. CHCl₃ acetylated overnight at room temperature with 1 ml. Ac₂O and 4 ml. C₅H₅N gave 87 mg. I diacetate (XV), m. 275-7°, [α]_D -13° (c 1.1, CHCl₃). XV (80 mg.) in 6 ml. alc. free CHCl₃ concentrated to 5 ml. and diluted with 4 ml. C₅H₅N, the mixture treated dropwise at -15° with 12 ml. 10% COCl₂-MePh and the mixture warmed at 15° 1 hr., kept overnight at 15-20° and the COCl₂ decomposed with ice, the mixture diluted with H₂O and Et₂O and the washed and dried Et₂O layer evaporated, the gum chromatographed on 2 g. Al₂O₃, and eluted with C₆H₆ gave 37 mg. 25D-spirostan-1 β ,2 β ,3 α ,5 β -tetrol 1,5-carbonate 2,3-diacetate, m. 169-72°, [α]_D 27° (c 1.1, CHCl₃), λ 5.66, 5.72, 8.08, 8.19, 8.40 μ (CS₂), no OH band. Further elution with 1:1 CHCl₃MeOH yielded 15 mg. impure XV, m. 257-66°. The likelihood that I should have the same cis configuration of the A/B ring junction as yonogenin and tokorogenin was strengthened by the close

resemblance of the rotatory dispersion curves of XII and 25D,5 β -spirostan-1 β ,2 β -diol-3-one acetonide, derived from tokorogenin acetonide. From the formation of XV it was concluded that the OH group at C-5 in I was β -oriented and accordingly I was described as 25D-spirostan-1 β ,2 β ,3 α ,5 β -tetrol.


IT 6869-44-9, 25D-Spirost-4-ene-1 β ,2 β ,3 α -triol
133326-87-1, Kogagenin, anhydro-
(and derivs.)
RN 6869-44-9 HCAPLUS
CN Spirost-4-ene-1,2,3-triol, (1 β ,2 β ,3 α ,25R)- (9CI) (CA
INDEX NAME)

Absolute stereochemistry.




RN 133326-87-1 HCAPLUS
CN Kogagenin, anhydro- (6CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 547-01-3P, Tokorogenin 1255-15-8P, 25D-Spirosta-1,4-dien-
3-one, 2-hydroxy-
RL: PREP (Preparation)
(preparation of)
RN 547-01-3 HCAPLUS
CN Spirostan-1,2,3-triol, (1 β ,2 β ,3 α ,5 β ,25R)- (9CI) (CA
INDEX NAME)


Absolute stereochemistry.



RN 1255-15-8 HCAPLUS

CN Spirosta-1,4-dien-3-one, 2-hydroxy-, (25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 9 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1958:88167 HCAPLUS [Full-text](#)

DN 52:88167

OREF 52:15559f-i,15560a-b

TI The structure of tokorogenin

AU Morita, Katsura

CS Takeda Pharm. Inds., Ltd., Osaka

SO Pharmaceutical Bulletin (1957), 5, 494-6

CODEN: PHBUA9; ISSN: 0369-9471

DT Journal

LA Unavailable

GI For diagram(s), see printed CA Issue.

AB The structure of the previously isolated (Nishikawa, et al., C.A. 49, 14785d) tokorogenin (I) is here established. Oxidation of I by CrO₃ in AcOH gave tokorogenic acid (II), m. 250°; [α]_D²³ -26.3°; anhydride (with Ac₂O) (III), m. 268° (v (Nujol) 1800, 1755 cm.⁻¹); di-Me ester (with CH₂N₂) (IV), m. 157°. II with MeOH-HCl gave the α-mono-Me ester (V), m. 185°, also formed from III with MeONa; whereas IV hydrolyzed by NaOH gave the β-mono-Me ester, m. 208°. I gave its acetonide (VI), m. 303°, hydrolyzed back to I by hot AcOH. VI with p-MeC₆H₄SO₂Cl in C₅H₅N gave its tosyl ester, m. 203° (decomposition), which was hydrolyzed by hot AcOH to the 3-tosyl ester of I, and this in turn gave with MeOH-KOH the epoxide (VII), m. 235°. VII (oxidized by CrO₃ in C₅H₅N gave the α,β-epoxyketone (VIII), m. 236°, which treated with CrCl₂ gave the α,β-unsatd. ketone, m. 219° (λ 225 mμ at ε 7690), catalytically hydrogenated (Pd-C) to the saturated ketone, m. 182°, and this was finally reduced by the Huang-Minlon reaction to the known 5β-25D-spirostan (IX), m. 137°. Thus the 1- and 2-HO groups in I are shown to be cis, and the 3-HO group trans.

Reduction of VII by LiAlH_4 gave $1\beta,3\beta$ -dihydroxysapogenin, m. 238° , which formed neither an acetonide nor an epoxy compound. A 2nd series of reactions led also to IX from VI. Oxidation of VI by CrO_3 in $\text{C}_5\text{H}_5\text{N}$ gave the 3-oxo derivative (HO changed to $:\text{O}$) (X), m. 229° , which was hydrolyzed by hot AcOH to the dihydroxyketone (XI), m. 225° . With alkali, both X and XI gave the enol form of the α -diketone (XII), m. 225° , (λ 269 $\text{m}\mu$ at ϵ 6900), oxidized by alkaline H_2O_2 to the known samogenic acid, m. 270° ; $[\alpha]_{23\text{D}} -37^\circ$; di-Me ester, m. 147° . Both X and XII with alkaline $\text{N}_2\text{H}_4 \cdot \text{H}_2\text{O}$ gave IX. From all these results I was established as $1\beta,2\beta,3\alpha$ -trihydroxy- 5β -25D-spirostan.


IT 547-01-3, Tokorogenin

(and cyclic 1,2-acetal with acetone and other derivs.)

RN 547-01-3 HCAPLUS

CN Spirostan-1,2,3-triol, ($1\beta,2\beta,3\alpha,5\beta,25\text{R}$)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



(and derivs., as structure for tokorogenin)

IT 472-10-6P, $5\beta,25\text{D}$ -Spirostan- $1\beta,3\beta$ -diol

6870-82-2P, $5\beta,25\text{D}$ -Spirostan-3-one, $1\beta,2\beta$ -dihydroxy-


RL: PREP (Preparation)

(preparation of)

RN 472-10-6 HCAPLUS

CN Spirostan-1,3-diol, ($1\beta,3\beta,5\beta,25\text{R}$)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 6870-82-2 HCAPLUS

CN 5β -Spirostan-3-one, $1\beta,2\beta$ -dihydroxy-, (25R)- (8CI) (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 10 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1958:65895 HCAPLUS Full-text

DN 52:65895

OREF 52:11878e-i,11879a-g

TI Reduction of $4\beta,5$ -epoxysmilagenone and $4\alpha,5$ -epoxytigogenone with lithium aluminum hydride

AU de Vivar, A. Romo; Ruelas, J. Perez; Romo, J.

SO Boletin del Instituto de Quimica de la Universidad Nacional Autonoma de Mexico (1957), 9, 59-72

CODEN: BIQUA5; ISSN: 0076-745X

DT Journal

LA Unavailable

OS CASREACT 52:65895

AB Oxidation of Δ^4 -diosgenone (I) with H_2O_2 in an alkaline medium yielded 2 isomeric epoxides, $4\beta,5$ -epoxysmilagenone (II), and $4\alpha,5$ -epoxytigogenone (III). Reduction of II and III with $LiAlH_4$ yielded 4 diols, 2 each from II and III. M.ps. were uncor., rotations determined at 23° in $CHCl_3$. 2α -Acetoxydiosgenone (4 g.) in 30 ml. tetrahydrofuran (THF) added slowly to 1 g. $LiAlH_4$ in 50 ml. Et₂O, the mixture refluxed 1 hr., poured into H₂O, acidified with dilute HCl, extracted with $CHCl_3$, and the washed and dried extract evaporated yielded 2.2 g. $20\alpha,22\alpha$ -25D-spirost-4-ene- $2\alpha,3\beta$ -diol (IV), m. $251-4^\circ$, $[\alpha]_D -83.5^\circ$; diacetate (V), m. $209-11^\circ$, $[\alpha]_D -136^\circ$. V (400 mg.) in 50 ml. EtOAc and 20 ml. AcOH hydrogenated over 80 mg. PtO₂ and the mixture filtered and concentrated yielded 260 mg. gitogenin, m. $271-3^\circ$, $[\alpha]_D -65^\circ$. I (10 g.) in 400 ml. EtOH treated simultaneously at room temperature with 3 g. KOH in 6 ml. H₂O and 15 ml. 30% H_2O_2 , the mixture held 45 min. at 30° , diluted with H₂O, and filtered yielded 3.45 g. II, m. $211-12^\circ$, $[\alpha]_D 20^\circ$. The mother liquors dissolved in hexane and chromatographed on Al_2O_3 yielded 350 mg. III, m. $210-11^\circ$, $[\alpha]_D -125^\circ$. II (1 g.) in 20 ml. AcOH treated with 2 ml. H_2SO_4 in 10 ml. AcOH, the mixture held 1 hr. at room temperature, diluted with H₂O, and filtered yielded 0.5 g. 4-hydroxydiosgenone (VI), m. $241-2^\circ$, $[\alpha]_D -30^\circ$; acetate, m. $203-4^\circ$ (Ac₂O-pyridine), $[\alpha]_D -13^\circ$. III (300 mg.) in 20 ml. AcOH treated with 1 ml. H_2SO_4 in 10 ml. AcOH yielded 125 mg. VI, m. $238-40^\circ$. VI (300 mg.) and 500 mg. o-C₆H₄(NH₂)₂ refluxed 2 hrs., diluted with H₂O, and filtered yielded 140 mg. phenazine derivative, m. $247-9^\circ$, $[\alpha]_D -61^\circ$. II (10 g.) in 70 ml. THF added slowly to 4 g. $LiAlH_4$ in 100 ml. Et₂O, the mixture refluxed 1 hr., decomposed with EtOH, poured into H₂O, acidified with dilute HCl, and filtered yielded 3.45 g. 5-hydroxyepismilagenin (VII), m. $260-3^\circ$, $[\alpha]_D -60^\circ$; 3-monoacetate (VIIa), m. $218-20^\circ$ (Ac₂O-pyridine), $[\alpha]_D -45^\circ$. The mother liquors from VII evaporated to dryness, the residue (6.2 g.) heated 1 hr. on the steam bath with 30 ml. Ac₂O and 30 ml. pyridine, diluted with H₂O,

and filtered yielded 1.7 g. 3-acetate (VIII), m. 203-5°, of 5-hydroxyepismilagenin. The mother liquor chromatographed on Al_2O_3 and eluted with hexane yielded 3.16 g. 3-monoacetate (IX), m. 195-6°, $[\alpha]_D -42^\circ$, of 5-hydroxysmilagenin. The end fractions yielded 440 mg. VIIa, m. 215-17°. IX (1 g.) in 100 ml. MeOH treated with 500 mg. K_2CO_3 in 8 ml. H_2O , the mixture refluxed 1 hr., diluted with H_2O , and filtered yielded 680 mg. 5-hydroxysmilagenin (X), m. 265-7°, $[\alpha]_D -46^\circ$. X (880 mg.) dissolved in 50 ml. CHCl_3 , 15 ml. CHCl_3 distilled, 3 ml. pyridine added, the mixture treated at 0° with 1 ml. SOCl_2 , held 3 hrs. at 0°, washed, and concentrated yielded 575 mg. cyclic sulfite (XI), m. 193-4°, $[\alpha]_D -74^\circ$. X (300 mg.) in 15 ml. AcOH treated with 150 mg. CrO_3 in 0.5 ml. H_2O and 4 ml. AcOH, the mixture allowed to stand 1 hr. at room temperature, poured into H_2O , and filtered yielded 80 mg. 5-hydroxysmilagenone (XIIa), m. 240-2°, $[\alpha]_D -32^\circ$. VII (2 g.) in 40 ml. AcOH containing 750 mg. CrO_3 yielded 1.2 g. XIIa, m. 240-2°. XIIa (300 mg.) in 30 ml. MeOH refluxed 1 hr. with 1 ml. HCl, diluted with H_2O , and filtered yielded 230 mg. I, m. 188-90°, $[\alpha]_D -19^\circ$. XIIa (400 mg.) in 30 ml. MeOH treated with 500 mg. KOH in 2 ml. H_2O and the mixture refluxed yielded 250 mg. I, m. 183-5°. IX (1 g.) and 100 mg. p-MeC₆H₄SO₃H in 25 ml. Ac₂O held overnight at room temperature, the mixture poured into H_2O , and filtered yielded 690 mg. diacetate (XIII), m. 208-10°, $[\alpha]_D -55^\circ$. XIII (500 mg.) in 50 ml. MeOH containing 500 mg. KOH refluxed 9 hrs., diluted with H_2O , and filtered yielded 380 mg. X, m. 248-52°. XIII (500 mg.) refluxed 1 hr. with 500 mg. K_2CO_3 was recovered. IX (1 g.) and 100 mg. p-MeC₆H₄SO₃H in 40 ml. Ac₂O processed as for VIII yielded 880 mg. diacetate (XIV), m. 248-50°, $[\alpha]_D -66^\circ$. XIV (1.96 g.) refluxed 1 hr. in 250 ml. MeOH containing 1.5 g. KOH, the solution concentrated to 1/2 the original volume, diluted with H_2O , and filtered yielded 1.585 g. 5-acetoxyepismilagenin (XV), m. 215-16°, $[\alpha]_D 80^\circ$. XV (1.35 g.) in 60 ml. AcOH at room temperature treated with 500 mg. CrO_3 in 10 ml. 80% AcOH, the mixture held 2 hrs. at room temperature, and processed as for XIIa yielded 930 mg. 5-acetoxysmilagenone (XVI), m. 213-15°, $[\alpha]_D -49^\circ$. XVI with HCl or KOH yielded I, m. 180-3° and 185-7°, resp. III (400 mg.) in 10 ml. Et₂O added slowly to 400 mg. LiAlH_4 in 20 ml. Et₂O, the mixture refluxed 30 min., poured into H_2O , acidified with dilute HCl, and extracted with CHCl_3 yielded 110 mg. 5-hydroxytigogenin (XVII), m. 265-7°, $[\alpha]_D -50^\circ$. XVII (1 g.) in 20 ml. oxidized with 350 mg. CrO_3 in 6 ml. 80% AcOH yielded 660 mg. 5-hydroxytigogenone (XVIII), m. 278-80°, $[\alpha]_D -60^\circ$. 5-Hydroxyepitigogenin (60 mg.) in 3 ml. AcOH treated with 30 mg. CrO_3 in 3 ml. 80% AcOH yielded 16 mg. XVIII, m. 273-5°. XVIII with HCl yielded I, m. 182-4°. The monoacetate (1 g.) treated with Ac₂O and p-MeC₆H₄SO₃H and the product chromatographed on Al_2O_3 yielded 510 mg. 5-hydroxytigogenin diacetate (XIX), m. 195-6°, $[\alpha]_D -64^\circ$. XIX (300 mg.) refluxed 1 hr. in 20 ml. MeOH containing 500 mg. KOH yielded 265 mg. 5-acetoxytigogenin (XX), m. 210-11°, $[\alpha]_D -62^\circ$. XX (250 mg.) in 10 ml. AcOH oxidized with 150 mg. CrO_3 in 5 ml. 80% AcOH yielded 135 mg. 5-acetoxytigogenone (XXI), m. 218-19°, $[\alpha]_D -78^\circ$. Dehydration with HCl yielded I, m. 183°. I (4 g.) in 70 ml. CHCl_3 treated with 4 g. (BzO)₂ in 100 ml. CHCl_3 , the mixture held 72 hrs. at 4°, washed, and the solvent evaporated yielded 2.1 g. 5,6 α -epoxytigogenin (XXII), m. 222-3°, $[\alpha]_D -118^\circ$; acetate, m. 236-7° (Ac₂O-pyridine 1 hr. on the steam bath), $[\alpha]_D -122^\circ$. XXII (1.3 g.) in 15 ml. THF added to 800 mg. LiAlH_4 in 60 ml. Et₂O, the mixture refluxed 3 hrs., poured into H_2O , acidified with dilute HCl, heated slightly to evaporate the Et₂O, then filtered yielded 900 mg. XVII, m. 264°; acetate, m. 242-3°.


IT 13944-32-6P, Diosgenone, 4-hydroxy-, phenazine derivative, acetate
119008-61-6P, 20 α ,22 α ,25D-Spirost-4-ene-2 α ,3 β -diol

RL: PREP (Preparation)
(preparation of)

RN 13944-32-6 HCAPLUS

CN Spirost-4-en-3-one, 4-hydroxy-, (25R)- (8CI, 9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 119008-61-6 HCAPLUS

CN 20 α , 22 α , 25D-Spirost-4-ene-2 α , 3 β -diol (6CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 262437-80-9, Aluminum lithium hydride
(reduction with, of 4 β , 5-epoxysmilagenone and
4 α , 5-epoxytigogenone)

RN 262437-80-9 HCAPLUS

CN Aluminum lithium hydride (CA INDEX NAME)

| Component | Ratio | Component Registry Number |
|-----------|-------|------------------------------|
| H | x | 12385-13-6 |
| Li | x | 7439-93-2 |
| Al | x | 7429-90-5 |

L97 ANSWER 11 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1955:77946 HCAPLUS Full-text

DN 49:77946

OREF 49:14785h-i, 14786a-f

TI Constitution and stereochemistry of samogenin, markogenin, and mexogenin

AU Djerassi, Carl; Fishman, Jack; Moore, James A.

CS Wayne Univ., Detroit, MI

SO Chemistry & Industry (London, United Kingdom) (1954) 1320-2


CODEN: CHINAG; ISSN: 0009-3068

DT Journal
LA Unavailable

AB Samogenin (I) was converted to the dimesylate (II), C₂₉H₄₈O₈S₂, m. 201-2° (all m.ps. uncorr.), [α]_{28D} -63° (all rotations in CHCl₃). II with NaI in Me₂CO gave an olefin (III), C₂₇H₂₂O₂, m. 149-50°, [α]_{26D} -84°. Reduction of III with Pt oxide in EtOH yielded 22a-spirostan (IV), C₂₇H₄₄O₂, m. 139-40°, [α]_{25D} -75°, also obtained by Wolff-Kishner reduction of 3-oxo-22a-spirostan, which in turn had been derived from diosgenin (C.A. 47, 8761d). This constitutes the 1st rigorous correlation of I with a known sapogenin and establishes the stereochemistry of all asymm. centers with the exception of the 2 vicinal OH groups. I readily forms an acetonide, C₃₀H₄₈O₄, m. 167-70°, [α]_{29D} -72°, under conditions where gitogenin (2α,3β-dihydroxy-5α,22a-spirostan) was recovered, indicating that I is a cis glycol. This was confirmed when III was treated with OsO₄; the resulting glycol, m. 205-7°, [α]_D -88°, (diacetate, m. 196-8°, [α]_D -75°), was identical with the natural sapogenin. This indicates that I is a cis glycol of the 5β ("normal") series with the OH groups located most probably at positions 2 and 3 or 3 and 4. A differentiation between these 2 alternatives was accomplished in the following manner. Wolff-Kishner reduction of 3-oxo-22a-spirostan-4-ene (diosgenone) gave 3 olefins, C₂₇H₄₂O₂, separable by chromatography: (a) 22a-spirostan-4-ene, m. 134-5° and 144-6°, [α]_{23D} -30°, also obtained by Raney Ni desulfurization of diosgenone cycloic ethylene mercaptal, C₂₉H₄₄O₂S₂, m. 265-7°, [α]_{25D} 30°; (b) 5α,22a-spirostan-3-ene, m. 172-4°, [α]_{25D} -34°, converted by catalytic hydrogenation to 5α,22a-spirostan and by BzO₂H oxidation to the corresponding 3α,4α-oxide, C₂₇H₄₂O₃, m. 195-8°, [α]_{22D} -60°, the structure of which was demonstrated by LiAlH₄ reduction to epitigogenin (3α-hydroxy-5α,22a-spirostan); and (c) an olefin which was assigned the structure 22a-spirostan-3-ene (V), m. 142.5-44°, [α]_{25D} -86°, since it was hydrogenated readily to IV. OsO₄ hydroxylation of V yielded the corresponding 3ξ,4ξ(cis)-dihydroxy-22a-spirostan (VI), C₂₇H₄₄O₄, m. 192-5°, [α]_{29D} -82° (diacetate, m. 210-12°, [α]_{29D} -46°), which was oxidized with CrO₃ to the derived 3,4-seco acid (VII), C₂₇H₄₂O₆, m. 264-6°, [α]_{27D} -19° (C₅H₅N) [di-Me ester (VIII), m. 195-7°, [α]_{25D} -47°]. VII and VIII were not identical to the corresponding oxidation products of I, samogenic acid, m. 270-3°, [α]_{27D} -39° (C₅H₅N), and di-Me samogenate (IX), m. 145-7°, [α]_{22D} -32°, thus excluding a 3,4-di-HO structure for I. 3α-Hydroxy-22a-spirostan tosylate, C₃₄H₅₀O₅S, m. 168-70°, [α]_{29D} -38°, yielded a mixture of the Δ²- and Δ³-olefins, III and V, when refluxed with collidine. OsO₄ hydroxylation of this product, followed by chromatographic separation, furnished I and VI. In a 2nd experiment, the hydroxylation product was not purified but rather converted by CrO₃ oxidation, CH₂N₂ methylation, and chromatography into VIII and IX. This reduces the structural possibilities for I to 2α,3α- and 2β,3β-dihydroxy-22a-spirostan with the same configuration in the side chain as diosgenin. Since markogenin (X) affords a pseudo derivative, different from that derived from I, but can be isomerized by strong acid to I, it follows that X differs from I only in the configuration at C-25 and possibly also at C-22. The same configuration for the HO groups has been demonstrated previously. Since mexogenin yields I on Wolff-Kishner reduction, it must be x-oxo-22a-spirostan-2α,3α- or 2β,3β-diol. This group of 2,3-dihydroxy-5β-sapogenins represents the 1st example of naturally occurring cis glycols in the sapogenin series.


IT 639-95-2, 5β,22a-Spirostan-3-one
(Wolff-Kishner reaction with)
RN 639-95-2 HCAPLUS
CN Spirostan-3-one, (5β)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 7662-01-3, 22a-Spirost-4-en-3-one
 (Wolff-Kishner reaction with, and cyclic ethylene mercaptole and other
 derivs.)
 RN 7662-01-3 HCAPLUS
 CN Spirost-4-en-3-one (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 911458-95-2, 5β,22a-Spirostan-2β,3β-diol
 911459-00-2, 5β,22a-Spirostan-2α,3α-diol
 (and derivs.)
 RN 911458-95-2 HCAPLUS
 CN 5β,22a-Spirostan-2β,3β-diol (5CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 911459-00-2 HCAPLUS
 CN 5β,22a-Spirostan-2α,3α-diol (5CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 6788-40-5P, Epitigogenin 61010-49-9P,
 22a-Spirostan-3 ξ , 4 ξ -diol 911453-64-0P,
 5 α , 22a-Spirostan-3 α -ol
 RL: PREP (Preparation)
 (preparation of)
 RN 6788-40-5 HCAPLUS
 CN Spirostan-3-ol, (3 α , 5 α , 25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 61010-49-9 HCAPLUS
 CN Spirostan-3, 4-diol, (3 α , 4 β , 5 α , 25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 911453-64-0 HCAPLUS

CN 5 α ,22a-Spirostan-3 α -ol (5CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 562-35-6, Markogenin 16680-64-1, Mexogenin
(stereochemistry of)

RN 562-35-6 HCAPLUS

CN Spirostan-2,3-diol, (2 β ,3 β ,5 β ,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 16680-64-1 HCAPLUS


CN Spirostan-12-one, 2,3-dihydroxy-, (2 β ,3 β ,5 β ,25R)- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



IT 469-97-6, Samogenin
 (stereochemistry of, and cyclic acetal with acetone and other derivs.)
 RN 469-97-6 HCAPLUS
 CN Spirostan-2,3-diol, (2 β ,3 β ,5 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L97 ANSWER 12 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN
 AN 1955:49607 HCAPLUS Full-text
 DN 49:49607
 OREF 49:9685g-i,9686a-i,9687a-g
 TI Synthesis of cortisone. VIII. Wagner-Meerwein rearrangement involving rings C and D of the steroid nucleus
 AU Elks, J.; Phillipps, G. H.; Taylor, D. A. H.; Wyman, L. J.
 CS Natl. Inst. Med. Research, London
 SO Journal of the Chemical Society (1954) 1739-49
 CODEN: JCSOA9; ISSN: 0368-1769
 DT Journal
 LA Unavailable
 AB cf. C.A. 49, 2470i. Hecogenin acetate (I) (100 g.) and 100 g. p-MeC₆H₄SO₂NHNH₂ (II), in 7 l. EtOH refluxed 65 hrs. gave 100 g. (74%) crude hecogenin acetate p-toluenesulfonylhydrazone (III); m. 274° (decomposition) (from EtOH-CHCl₃), [α]_D -15° (all rotations determined in CHCl₃ unless otherwise stated), λ_{EtOH}max. 226 mμ (ε 11600), ν_{maximum} (CS₂) 3250, 1732, 1245, 1342, 1160, 977, 914, 897, and 860 cm.⁻¹ In another experiment with 36 g. I, 18.6, 24.9, and 31.1 g. III were deposited after 6, 24, and 50 hrs. I (20 g.) in CHCl₃ added to 10 g. II in EtOH and concentrated HCl gave 23 g. (85%) III. I (2 g.) in HOAc treated 1 hr. with 2 g. II yielded 2.4 g. (89%) III. III (50 g.) heated gradually to 130-40° with 15 g. Na in 1 l. (CH₂OH)₂ until no further gas evolution was apparent, and the mixture cooled, diluted with H₂O, and left overnight yielded on crystallization 21.4 g. of a free alc. (IV), m. 125-33°. IV on acetylation yielded 19.5 g. (55%) "compound A" (V), m. 142-5° (from aqueous MeOH), [α]_D -57°, ν_{maximum} (CS₂) 1733, 1238, 980, 920, 898 and 860 cm.⁻¹ Purified IV, obtained by hydrolysis of V with alc. KOH, m. 120-5°, [α]_D -55°, ν_{maximum} (Nujol) 3400, 983, 920, 898 and 865 cm.⁻¹ The mother liquor from III yielded 1.2 g. solid which on acetylation yielded 0.94 g. 3 β -acetoxy-5 α ,22a-spirost-11-ene (VI), m. 206-11°. An alternative method of isolating V was by acetylating the crude reaction product, followed by chromatography. III (28 g.) refluxed 0.5 hr. with 3 g. Na in 800 ml. BuOH and the crude product fractionally crystallized yielded 9.18 g. (51%) IV and 2.1 g. (12%) 5 α ,22a-spirost-11-en-3 β -ol (VII), tablets, m. 192-4°, [α]_D -37°. VII on acetylation yielded VI, [α]_D -43°. Acetylation of the solid obtained from the combined mother liquors gave 2.6 g. VI (total yield, 25%). The mother

liquor yielded a solid which on chromatography gave 700 mg. (3.5%) 3 β -acetoxy-C-nor-D-homo-5 α ,22a-spirost-17a-ene (VIII), m. 215°, [α]_D -81°. V (0.912 g.) in CCl₄ treated at -25° with 320 mg. Br in CCl₄, and the solution warmed up to 0°, then washed with H₂O, NaHCO₃ solution, and H₂O, yielded 0.81 g. (66%) of the dibromide (IX), m. 108° (decomposition), [α]_D -33°, ν _{max} (CS₂) 1735, 1240, 988, 918, 898, 860, and 702 cm.⁻¹ IX decomposed when kept at room temperature or on attempted crystallization Tigogenin acetate (X) (0.5 g.) in HOAc hydrogenated at room temperature and pressure with 200 mg. PtO₂ took up 1 mole H in 1 hr. and the residue on acetylation yielded 0.11 g. X, m. 204-7°, [α]_D -72°, and 0.2 g. 3 β ,26-diacetoxy-5 α ,22a-furostan (XI), m. 114-16°, [α]_D -14°, ν _{max} (CS₂) 1732 and 1240 cm.⁻¹ The bands characteristic of the 22a-spirostan system were almost absent. XI hydrolyzed with alkaline KOH yielded 3 β ,26-dihydroxy-5 α ,22a-furostan, m. 165-7°, [α]_D -6°, ν _{max} (Nujol) 3300 cm.⁻¹ V (5 g.) in CHCl₃ left 0.5 hr. with 15 ml. 2.8N o-HO₂CC₆H₄CO₃H in Et₂O at room temperature and the crude product chromatographed on Al₂O₃ yielded 1.06 g. (21%) epoxide P (XII), m. 189-90°, [α]_D -66°, ν _{max} (CS₂) 1732, 1240, 980, 918, 895, and 860 cm.⁻¹, and 2.01 g. (39%) epoxide Q (XIII), m. 194-5°, [α]_D -63°, ν _{max} (CS₂) 1734, 1240, 980, 918, 895, and 860 cm.⁻¹ The infrared spectra of XII and XIII are quite different in detail. XII (0.5 g.) in 50 ml. tetrahydrofuran refluxed 3.5 hrs. with 0.5 g. LiAlH₄ and the crude product acetylated yielded 41 mg. starting material and 0.383 mg. (76%) diol monoacetate (XIV), prisms, m. 192-4°, [α]_D -63°, ν _{max} (CS₂) 3620, 1732, 1240, 978, 918, 898, and 860 cm.⁻¹ XIV (100 mg.) left 2 hrs. at room temperature in 2 ml. C₅H₅N with 0.5 ml. POCl₃ yielded 51 mg. V. XIII (0.5 g.) similarly reduced with LiAlH₄ 5 hrs. and the crude product acetylated yielded 0.306 g. (61%) of a diol monoacetate (XV), prisms, m. 161°, resolidified in needles, and finally m. 170-1°, [α]_D -53° (c 1.4, Me₂CO), ν _{max} (CS₂) 3600, 1735, 1240, 980, 916, 897, and 860 cm.⁻¹ XV (0.155 g.) in 2 ml. C₅H₅N similarly dehydrated with POCl₃ gave 0.092 g. V. V (1.6 g.) in Et₂O containing 0.7 C₅H₅N left 65 hrs. with 1 g. OsO₄, the Et₂O removed, the residue refluxed 4.5 hrs. with 7 g. Na₂SO₃ in EtOH and H₂O, and the crude product acetylated yielded 1.18 g. (69%) of a triol monoacetate (XVI), m. 214-17°, [α]_D -40°, ν _{max} (CS₂) 3620, 1732, 1240, 978, 918, 896, and 860 cm.⁻¹ XVI was hydrolyzed to the free triol (XVII), prisms, m. 229-33°, [α]_D -40°, ν _{max} (CS₂) 3350, 982, 918, 900, and 860 cm.⁻¹ XVII (0.46 g.) in MeOH left 2 days at room temperature with 5 ml. 10% aqueous HIO₄ yielded 0.28 g. (61%) of a diketone (XVIII), m. 157-60°, [α]_D -21°, ν _{max} (CS₂) 3620, 1740, and 1714 cm.⁻¹ XVIII gave a violet color with Na nitroprusside, showing the presence of a Me ketone grouping. I (10 g.) in EtOH and CH₂Cl₂ left 3 days at room temperature with 0.6 g. NaBH₄ in 5 ml. H₂O gave 4.2 g. (42%) 3 β -acetoxy-5 α ,22a-spirostan-12 β -ol (XIX), m. 211-16°, contaminated with the 12 α -HO isomer. XIX (4 g. crude) added to 3 ml. MeSO₂Cl in 13 ml. C₅H₅N at 0°, left overnight at room temperature, the crude product refluxed 2 hrs. with 100 ml. MeOH, the solution evaporated to dryness, and the residue hydrolyzed by refluxing 0.5 hr. with 4% KOH in 80% EtOH yielded 1.03 g. IV. The mother liquor from IV evaporated to dryness, and the residue reacetylated and chromatographed gave 800 mg. V and 1.26 g. 3 β -acetoxy-12 α -methanesulfonyloxy-5 α ,22a-spirostan (XX), m. 186-8°, [α]_D -21°. The mother liquor from XX gave upon chromatography of the solid 75 mg. VIII. I (5 g.) in 25 ml. tetrahydrofuran refluxed 1 hr. with 0.8 g. LiAlH₄ in 25 ml. tetrahydrofuran and the crude product acetylated yielded after chromatography 10% 3 β ,12 α -diacetoxy-5 α ,22a-spirostan (XXI), prisms from aqueous MeOH, m. 153-6°, [α]_D -17° (Me₂CO, c 1), ν _{max} (CS₂) 1738, 1240, 976, 918, 895, and 860 cm.⁻¹ XXI was hydrolyzed with EtOH-KOH to the free diol (XXII), m. 200-6°, [α]_D -30° (Me₂CO), ν _{max} (Nujol) 3600, 3450, 981, 919, 903, and 863 cm.⁻¹ The chromatogram also yielded 45% 3 β ,12 β -diacetoxy-5 α ,22a-spirostan (rockogenin

diacetate) (XXIII), m. 198-203°, $[\alpha]_D$ -68° (c 1, CHCl₃), -63° (Me₂CO), ν_{maximum} (CS₂) 1735, 1240, 978, 918, 898, and 860 cm.⁻¹ XXIII similarly yielded the free diol (XXIV), m. 216-19°, $[\alpha]_D$ -60° (Me₂CO), ν_{maximum} (Nujol) 3300, 973, 914, 892, and 860 cm.⁻¹ XXI (0.75 g.) refluxed 2 hrs. with 0.225 g. KHCO₃ in 24 ml. MeOH and 6 ml. H₂O gave 585 mg. (85%) crude 12 α -acetoxy-5 α ,22a-spirostan-3 β -ol (XXV) (pure, m. 231-3°), $[\alpha]_D$ -15°, ν_{maximum} (CS₂) 3620, 1739, 1240, 978, 920, 898, and 860 cm.⁻¹ XXV (0.55 g.) in 15 ml. HOAc left at room temperature 4 hrs. with 1.5 equivs. CrO₃ gave 0.5 g. (91%) 12 α -acetoxy-5 α ,22a-spirostan-3-one (XXVI), m. 214-17°, $[\alpha]_D$ 1°, ν_{maximum} (CS₂) 1739, 1237, 1715, 981, 920, 899, and 863 cm.⁻¹ XXVI was saponified with MeOH-KOH to 12 α -hydroxy-5 α ,22a-spirostan-3-one, m. 254-7°, $[\alpha]_D$ -30°, ν_{maximum} (CS₂) 3620, 1712, 979, 917, and 895 cm.⁻¹ IV (1 g.) in HOAc left at room temperature 4 hrs. with 17 ml. 0.55N CrO₃ yielded a ketone, m. 101-4°, $[\alpha]_D$ -40°, ν_{maximum} (CS₂) 1715, 980, 920, 900, and 860 cm.⁻¹; 2,4-dinitrophenylhydrazone, orange solid, m. 206-8°. 3 β -Acetoxy-12 α ,23-dibromo-5 α ,22a-spirostan-11 β -ol (3 g.) refluxed 3.5 hrs. with 30 g. Zn in 300 ml. HOAc yielded 1.68 g. (78%) VI, ν_{maximum} (CS₂) 1732, 1240, 978, 918, 895, and 860 cm.⁻¹, also obtained in 30% yield from 3 β -acetoxy-23-bromo-11 β ,12 β -epoxy-5 α ,22a-spirostan. The free alc., VII, obtained by hydrolysis of VI, ν_{maximum} (Nujol) 3560, 3330, 978, 918, 900, and 861 cm.⁻¹ VII like VI showed bands of medium intensity at 702 and 760 cm.⁻¹, with a shoulder at 3000 cm.⁻¹, indicative of a cis-1,2-disubstituted ethylene grouping. VI (200 mg.) in 6 ml. CHCl₃ left overnight in the refrigerator with 0.5 ml. 2.8N o-HO₂CC₆H₄CO₃H yielded 155 mg. (75%) 3 β -acetoxy-11 α ,12 α -epoxy-5 α ,22a-spirostan (XXVII), needles, m. 221-5°, $[\alpha]_D$ -49.5°, ν_{maximum} (CS₂) 1735, 1240, 978, 918, 895, and 860 cm.⁻¹ There was no indication of the presence of the isomeric 11 β ,12 β -epoxide. XXVII (0.5 g.) refluxed 2 hrs. with 0.6 g. LiAlH₄ in 25 ml. tetrahydrofuran yielded XXII. Crude XXII was acetylated to XXI, identical with the specimen prepared from I. XXI was obtained in 44% yield from XXVII. XXII (200 mg., crude) in 7 ml. HOAc left at room temperature 4 hrs. with CrO₃ gave hecogenone (XXVIII), plates, m. 232-5°, $[\alpha]_D$ 21°, ν_{maximum} (CS₂) 1710, 978, 918, 896, and 860 cm.⁻¹, identical with a sample prepared from rockogenin by the same method. 3 β -Acetoxy-5 α ,22a-spirostan-12 β -ol (rockogenin monoacetate) (XXIX), prepared by a method mentioned above, m. 214-19°, $[\alpha]_D$ -65° (CHCl₃), -61° (dioxane), ν_{maximum} (CS₂) 3620, 1736, 1238, 978, 918, 895, and 862 cm.⁻¹ XXIX with MeSO₂Cl yielded a solid, m. 125-30° (decomposition), which, refluxed 4 hrs. with 1.5 g. K in 100 ml. tert-BuOH, gave 2.8 g. (73%) 3 β -hydroxy-C-nor-D-homo-5 α ,22a-spirost-17-ene (XXX), m. 157-69°, $[\alpha]_D$ -66.5°, ν_{maximum} 1642, 886, 980, 920, 898, and 864 cm.⁻¹ The presence of an OH group was shown by the spectrum of a Nujol mull, with maximum at 3500 and 3280 cm.⁻¹ XXX yielded VIII, ν_{maximum} 1732, 1238, 1638, 884, 978, 918, 896, and 862 cm.⁻¹ The structure of compound A (V) is discussed in the light of both its reactions and the stereochem. requirements of the rearrangement.


IT 884309-78-8, 5 α ,22a-Spirostan-3 β ,12 α -diol
(and esters)
RN 884309-78-8 HCAPLUS
CN 5 α ,22a-Spirostan-3 β ,12 α -diol (5CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 863-85-4P, Rockogenin, acetate 2137-20-4P, Hecogenone
 16653-52-4P, Rockogenin 119065-01-9P,
 5 α ,22a-Spirost-11-en-3 β -ol 120964-63-8P,
 5 α ,22a-Spirost-11-en-3 β -ol, acetate 884310-29-6P,
 5 α ,22a-Spirostan-3 β ,12 β -diol 911442-63-2P,
 Methanesulfonic acid, 12-ester with 5 α ,22a-spirostan-
 3 β ,12 α -diol 3-acetate 911460-03-2P,
 5 α ,22a-Spirostan-3-one, 12 α -hydroxy-, acetate
 911460-08-7P, 5 α ,22a-Spirostan-3-one, 12 α -hydroxy-
 RL: PREP (Preparation)
 (preparation of)
 RN 863-85-4 HCAPLUS
 CN Spirostan-3,12-diol, 3-acetate, (3 β ,5 α ,12 β ,25R)- (9CI)
 (CA INDEX NAME)

Absolute stereochemistry.



RN 2137-20-4 HCAPLUS
 CN Spirostan-3,12-dione, (5 α ,25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 16653-52-4 HCAPLUS

CN Spirostan-3,12-diol, (3 β ,5 α ,12 β ,25R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 119065-01-9 HCAPLUS

CN 5 α ,22 α -Spirost-11-en-3 β -ol (6CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 120964-63-8 HCAPLUS

CN 5 α ,22a-Spirost-11-en-3 β -ol, acetate (6CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 884310-29-6 HCAPLUS

CN 5α,22a-Spirostan-3β,12β-diol (5CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 911442-63-2 HCAPLUS

CN Methanesulfonic acid, 12-ester with 5α,22a-spirostan-3β,12α-diol 3-acetate (5CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 911460-03-2 HCAPLUS

CN 5α,22a-Spirostan-3-one, 12α-hydroxy-, acetate (5CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 911460-08-7 HCAPLUS

CN 5α,22a-Spirostan-3-one, 12α-hydroxy- (5CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 915-35-5, Hecogenin, acetate
(spectrum of)

RN 915-35-5 HCAPLUS

CN Spirostan-12-one, 3-(acetyloxy)-, (3β,5α,25R)- (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 13 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1955:8351 HCAPLUS Full-text


DN 49:8351

OREF 49:1758g-i,1759a-e

TI Steroids. L. The oxidation of steroidal allylic alcohols with manganese dioxide. A novel synthesis of testosterone


AU Sondheimer, Franz; Amendolla, C.; Rosenkranz, G.
CS Syntex, S.A., Mexico City, Mex.
SO Journal of the American Chemical Society (1953), 75, 5930-2
CODEN: JACSAT; ISSN: 0002-7863
DT Journal
LA Unavailable
OS CASREACT 49:8351
AB cf. C.A. 48, 12157a. The oxidation of a number of steroidal allylic alcs. to the corresponding CO compds. with MnO₂ is described. 4-Androstene-3,17- dione (I) reduced with LiAlH₄ gave a mixture of 4-androstene-3 β ,17 β -diol (II) and the 3 α ,17 β -diol (III) which was oxidized with MnO₂ in 90% over-all yield to testosterone (IV). Similarly progesterone (V) was converted to 4-pregnen-20 β -ol-3- one (VI). The MnO₂ used in the oxidns. described was prepared from KMnO₄ and MnSO₄ as previously described (C.A. 48, 6386b). 4-Cholesten-3 β - ol, m. 129-31°, [α]D₂₀ 45°, in 100 cc. CHCl₃ shaken 24 h. at room temperature with 10 g. MnO₂ showed that the maximum at 240 m μ remained essentially unchanged and that an addnl. maximum at 284 m μ (log ϵ 3.17) appeared. A similar run shaken 3 h., and the resulting product recrystd. from MeOH yielded 0.93 g. (93%) 4-cholesten-3-one, m. 78-9°, λ maximum 240 m μ (log ϵ 4.22). A mixture of 22a-spirost-4-en-3 β -ol and the Δ 4-3 α -ol, m. 181-3°, obtained by the reduction of 22a-spirost-4-en-3-one, which with LiAlH₄ shaken in 150 cc. CHCl₃ with 15 cc. MnO₂ 4 h. at room temperature, the mixture filtered, and the product recrystd. from CHCl₃-Et₂O yielded 1.26 g. (84%) 22a-spirost-4-en-3-one, m. 183-5°, [α]D₂₀ -6, λ maximum 240 m μ (log ϵ 4.24); the rotations were measured in CHCl₃ and the ultra-violet absorption spectra in 95% EtOH. 22a-Spirost-5-en-3 β ,7 α -diol 3-acetate (0.50 g.), m. 190-3°, [α]D₂₀ -155°, in 50 cc. C₆H₆ shaken 24 h. at room temperature with 5 g. MnO₂, and the crystalline product [λ maximum 234 m μ (log ϵ 4.10)] recrystd. from MeOH yielded 0.29 g. (58%) 22a-spirost-5-en-3 β -ol-7-one acetate, m. 198-9°, [α]D₂₀ -158°, λ maximum 234 m μ (log ϵ 4.18), ν max. 1726, 1674 cm.⁻¹ 5 α ,22a-Spirost-9(11)-ene-3 β ,12-diol, m. 200-3°, (most probably a mixture of C-12 stereoisomers) in 50 cc. CHCl₃ shaken 10 h. at room temperature with 5 g. MnO₂, and the product [λ maximum 238 m μ (log ϵ 4.07)] recrystd. from CHCl₃-Me₂CO yielded 0.38 g. (76%) 5 α ,22a-spirost-9(11)-en-3 β -ol-12-one, m. 221-3°, λ maximum 238 m μ (log ϵ 4.16), ν max. 1718, 1670 cm.⁻¹ I (50 g.) in 300 cc. dry tetra-hydrofuran added with stirring and cooling to 15 g. LiAlH₄ in 1.5 l. THF during 0.5 h., the excess LiAlH₄ destroyed with EtOAc and concentrated aqueous Na₂SO₄, the mixture treated with 100 g. solid Na₂SO₄ and filtered, the filter residue washed with THF, and the solution evaporated yielded 50.4 g. mixture of II and III, m. 165-71°. The mixture ground in a mortar, suspended in 1250 cc. CHCl₃, stirred 10 h. at room temperature with 250 g. MnO₂, and filtered, the filter residue washed with hot CHCl₃, the combined filtrate and washing evaporated to dryness, and the residue recrystd. from Me₂CO-hexane yielded 38.2 g. IV, m. 152-3°, [α]D₂₀ 108°, λ maximum 240 m μ (log ϵ 4.23), 6.9 g. 2nd crop, m. 150-2°, and 3rd crops totaling 45.1 g. (90%). V (5.0 g.) reduced in the usual manner with LiAlH₄, the reduction product (5.0 g.), m. 162-72°, in 500 cc. CHCl₃ stirred 24 h. at room temperature with 50 g. MnO₂, and the product recrystd. from Et₂O-pentane gave 3.3 g. (66%) VI, m. 166-8°; recrystd., m. 174-5°, [α]D₂₀ 86°, λ maximum 240 m μ (log ϵ 4.23), λ max.CHCl₃ 1660 cm.⁻¹; acetate, m. 161-2° (from Me₂CO-hexane), [α]D₂₀ 134°, λ maximum 240 m μ (log ϵ 4.22), ν max. 1718, 1660 cm.⁻¹
IT 7662-01-3P, 22a-Spirost-4-en-3-one 882741-52-8P, 5 α ,22a-Spirost-9(11)-en-12-one, 3 β -hydroxy-
RL: PREP (Preparation)
(preparation of)
RN 7662-01-3 HCAPLUS
CN Spirost-4-en-3-one (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 882741-52-8 HCAPLUS
CN INDEX NAME NOT YET ASSIGNED

Relative stereochemistry.




L97 ANSWER 14 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN
AN 1954:3641 HCAPLUS Full-text
DN 48:3641
OREF 48:699f-i,700a-b
TI Steroids. XL. The oxidation of unsaturated steroidal alcohols with manganese dioxide
AU Sondheimer, F.; Rosenkranz, G.
CS Syntex, S. A., Laguna Mayran 413, Mexico City
SO Experientia (1953), 9, 62-3
CODEN: EXPEAM; ISSN: 0014-4754
DT Journal
LA English
OS CASREACT 48:3641
GI For diagram(s), see printed CA Issue.
AB cf. C.A. 47, 12415g. Vigorous shaking of Δ^4 -cholesten-3 β -ol or Δ^4 -22a-spirosten-3 β -ol with freshly precipitated MnO₂ resulted in conversion in satisfactory yield to the corresponding Δ^4 -3-ones in about 2 h. at room temperature. Similar oxidation of Δ^5 -22a-spirostene-3 β ,7 α -diol 3-acetate produced the Δ^5 -7-one; Δ^9 (11)-22a-5 α -spirostene-3 β ,12-diol gave the Δ^9 (11)-12-one; and Δ^5 ,17(20)-pregnadiene-3 β ,21-diol gave the corresponding 21-aldehyde, all in satisfactory yield. Δ^4 -cholestene-3 β ,6 β -diol in C₆H₆ with MnO₂ at room temperature was oxidized only at C-3 producing Δ^4 -cholesten-3-one-6 β -ol at 75% yield. This procedure has been used also to produce 6 β -hydroxyprogesterone

(m. 181-3°, $[\alpha]_{20D}$ 105°, all rotations in CHCl_3 , λ_{EtOHmax} . 236 $\text{m}\mu$, $\log \epsilon$ 4.22) and 6 β -hydroxy- Δ^4 -androstene-3,17-dione (m. 192-4°, $[\alpha]_{20D}$ 114°, λ_{EtOHmax} . 236 $\text{m}\mu$, $\log \epsilon$ 4.25). At reflux temperature this reaction produced the corresponding diketones. Δ^4 -androstene-3,17-dione was reduced with LiAlH_4 to a presumed mixture of Δ^4 -androstene-3 β ,17 β -diol and the 3 α ,17 β -diol which in CHCl_3 with MnO_2 at room temperature was only oxidized at C-3 to yield pure testosterone in 90% overall yield. The readily available Δ^5 -3 β -ols (Type I) with MnO_2 in refluxing C_6H_6 were found to yield the corresponding Δ^4 ,6-dien-3-ones (Type III) in conversions of about 30%. In this way the following dienones (Type III) were prepd: Δ^4 ,6-22a-spirostadien-3-one, Δ^4 ,6-cholestadien-3-one, Δ^4 ,6-androstadiene-3,17-dione, Δ^4 ,6-androstadien-17 β -ol-3-one (6-dehydrotestosterone), Δ^4 ,6-pregnadiene-3,20-dione (6-dehydroprogesterone), Δ^4 ,6-pregnadien-20 β -ol-3-one (m. 197-9° $[\alpha]_{20D}$ 15°, λ_{EtOHmax} . 282 $\text{m}\mu$, $\log \epsilon$ 4.54), Δ^4 ,6,16-pregnatriene-3,20-dione (from Δ^5 ,16-pregnadiene-3 β ,20 β -diol) (m. 253-6°, $[\alpha]_{20D}$ 144°, λ_{EtOHmax} . 240 and 284 $\text{m}\mu$, $\log \epsilon$ 4.21 and 4.53), Δ^4 ,6-pregnadien-17 α -ol-3,20-dione (m. 240-2°, $[\alpha]_{20D}$ 21°, λ_{EtOHmax} . 284 $\text{m}\mu$, $\log \epsilon$ 4.53), Δ^4 ,6 pregnadien-21-ol-3,20-dione acetate, and Δ^4 ,6-pregnadiene-17 α ,21-diol-3,20-dione 21-acetate (6-dehydro Reichstein's Substance S acetate) (m. 218-20°, $[\alpha]_{20D}$ 104°, λ_{EtOHmax} . 284 $\text{m}\mu$, $\log \epsilon$ 4.48). The reactions appear to pass through intermediates such as II.


IT 911461-28-4, 5 α ,22a-Spirost-9(11)-ene-3 β ,12-diol
(oxidation with manganese dioxide)
RN 911461-28-4 HCAPLUS
CN 5 α ,22a-Spirost-9(11)-ene-3 β ,12-diol (5CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 7662-01-3P, 22a-Spirost-4-en-3-one 37147-71-0P,
22a-Spirosta-4,6-dien-3-one 882741-52-8P, 5 α ,22a-Spirost-
9(11)-en-12-one, 3 β -hydroxy-
RL: PREP (Preparation)
(preparation of)
RN 7662-01-3 HCAPLUS
CN Spirost-4-en-3-one (9CI) (CA INDEX NAME)


Absolute stereochemistry.



Absolute stereochemistry.



Relative stereochemistry.



L97 ANSWER 15 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN
AN 1953:72869 HCAPLUS Full-text
DN 47:72869
OREF 47:12412c-i,12413a-e
TI The transformation of manogenin to hecogenin
AU Wendler, N. L.; Slates, H. L.; Tishler, M.
CS Merck & Co., Inc., Rahway, NJ
SO Journal of the American Chemical Society (1952), 74, 4894-7
CODEN: JACSAT; ISSN: 0002-7863
DT Journal

LA Unavailable

OS CASREACT 47:72869

GI For diagram(s), see printed CA Issue.

AB To 37.5 g. crude manogenin (I) containing 40-50% Δ^9 -dehydro derivative and varying amts. of gitogenin in 3 l. refluxing BuOH was added 75 g. Na portionwise as rapidly as possible, part of the BuOH removed in vacuo after 1-2 h. and the remainder as an azeotrope with H₂O, H₂O added to the residue, and the solid filtered and washed alkali-free to yield 35.6 g. agavogenin (II), feathery needles, m. 240-2° (from CHCl₃EtOAc). II refluxed 1 h. with Ac₂O gave the triacetate, m. 221-7° (from MeOH). II (35.5 g.) in 300 cc. dry pyridine heated 3 h. at 100°, under N, with 48 g. succinic anhydride, the mixture cooled, the pyridine removed in vacuo, the residue shaken with H₂O and CHCl₃, the aqueous layer extracted 3 times with CHCl₃, and the combined CHCl₃ extract washed with 2.5N HCl, H₂O, and saturated aqueous NaCl, dried and evaporated in vacuo, yielded 53 g. II bis(hemisuccinate) (III), not further purified. Crude III (53 g.) in 500 cc. AcOH oxidized at room temperature with 5.87 g. CrO₃ in 30 cc. 30% aqueous AcOH, the excess CrO₃ destroyed with MeOH, the solution concentrated to a small volume in vacuo, diluted with H₂O, extracted with CHCl₃-Et₂O, the extract washed with dilute H₂SO₄ and saturated aqueous NaCl, dried, and evaporated in vacuo, the residue (47 g.) dissolved in 1 l. MeOH containing 100 cc. H₂O and 100 g. KOH, the solution refluxed 4 h., the MeOH removed in vacuo, H₂O added, and the product extracted with CHCl₃ yielded 20.5 g. I, m. 254-7° (from CHCl₃EtOAc), containing some gitogenin. I (20 g.) in 200 cc. pyridine let stand 16 h. at 0-5° with 20 cc. MeSO₂Cl and the mixture poured with stirring into ice water gave 21.7 g. I dimesylate (methanesulfonate) (IV), long slender needles, m. 241° (decomposition) (from Me₂CO), $[\alpha]_D^{24.5}$ -44.2° (CHCl₃). IV (6.1 g.) heated 24 h. at 100° in a glass-lined autoclave with 15.25 g. NaI in 250 cc. dry Me₂CO, the mixture filtered, the residue washed with Et₂O and CHCl₃, the combined filtrate and washings concentrated in vacuo, and the residue diluted with CHCl₃ and Et₂O, washed with 5% aqueous Na₂S₂O₃ and H₂O, dried, and evaporated gave 3.75 g. crystals, m. 173-5°; 2.00 g. of the crystalline product in petr. ether chromatographed on acid-washed Al₂O₃ gave 800 mg. Δ^2 -isoallospirosten-12-one (V), micalike plates, m. 199-200° (from aqueous Me₂CO), $[\alpha]_D^{24.5}$ 39.1° (CHCl₃). Similarly hecogenin mesylate, m. 178°, was prepared and converted with NaI in Me₂CO, at 100° for 24 h., to V. From the mother liquor, of V, was obtained 950 mg. Δ^2 -22-isoallospirostene (VI), needles, m. 186-7° (from Me₂CO), giving a yellow color with C(NO₂)₄. Hecogenone (VII) (500 mg.), 6.0 g. KOH, 60 cc. (CH₂OH)₂, and 0.6 cc. 85% N₂H₄.H₂O heated cautiously to 140°, then 1 h. at 140°, and 1 h. at 190-5° while a slow stream of N was passed over the surface, poured into H₂O, and the precipitate washed alkali-free yielded 350 mg. 22-isoallospirostan-12-one (VIII), plates, m. 173-4°, $[\alpha]_D^{24.5}$ -61.8° (CHCl₃); also obtained by hydrogenation of VI in EtOAc over PtO₂. To 1 g. V in 15 cc. C₆H₆ was added, at 5°, 5 cc. C₆H₆ containing 0.3 g. BzO₂H, the mixture diluted with 100 cc. Et₂O, washed with cold 5% aqueous Na₂CO₃ and H₂O, and the Et₂O solution dried and evaporated to give 1.1 g. crude 2(α),3(α)-epoxy-22-isoallospirostan-12-one (IX), chromatographed on basic Al₂O₃ and recrystd. twice from Et₂O, m. 210-13°, $[\alpha]_D^{25}$ 22° (CHCl₃). To 400 mg. LiAlH₄ in 100 cc. dry Et₂O was added with vigorous stirring 1.1 g. IX in 20 cc. C₆H₆ and 40 cc. Et₂O, the mixture stirred 45 min. at room temperature, then refluxed 10 min., the excess hydride decomposed with H₂O and dilute HCl, the aqueous layer extracted with Et₂O, and the combined Et₂O layer and extract were washed acid-free with H₂O and saturated aqueous NaCl, dried, and evaporated to give 1.05 g. crude compound (X) in 20 cc. AcOH oxidized overnight at room temperature with 358 mg. CrO₃ in 15 cc. 80% AcOH gave 350 mg. VII, m. 238-41°, $[\alpha]_D^{24.5}$ 23.8° (CHCl₃); an addnl. 200 mg. VII was obtained from the mother liquor. VII (1.7 g.) in 75 cc. dry THF reduced with 1.5 g. LiAlH₄ gave crude compound (XI), which was dissolved in 20 cc. dry pyridine containing 3.0 g. succinic anhydride heated 3 h. on a steam bath under N, the mixture concentrated in


vacuo, diluted with H₂O, extracted with CHCl₃ and Et₂O, and the extract washed with dilute HCl, H₂O, and saturated aqueous NaCl, dried, and evaporated in vacuo to give 2.4 g. crude XI 3-hemisuccinate (XII). XII (2.4 g.) in 50 cc. AcOH oxidized 16 h. at room temperature with 350 mg. CrO₃ in 10 cc. 80% aqueous AcOH gave 2.16 g. crude hecogenin (XIII) hemisuccinate, which refluxed 4 h. under N with 75 cc. MeOH containing 4.0 g. KOH yielded 650 mg. XIII, small plates, m. 263-6°, [α]_D^{24.5} 13.5° (CHCl₃); acetate, m. 247-50° (from CHCl₃-EtOAc), [α]_D^{24.5} 92° (CHCl₃).

IT 467-55-0, Hecogenin
(and esters)

RN 467-55-0 HCAPLUS

CN Spirostan-12-one, 3-hydroxy-, (3 β , 5 α , 25R)- (CA INDEX NAME)

Absolute stereochemistry.




IT 511-96-6P, Gitogenin 564-43-2P, Manogenin
2137-20-4P, Hecogenone 883721-07-1P,
22-Isoallospirostan-3 α , 12-diol
RL: PREP (Preparation)
(preparation of)

RN 511-96-6 HCAPLUS

CN Spirostan-2,3-diol, (2 α , 3 β , 5 α , 25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 564-43-2 HCAPLUS

CN Spirostan-12-one, 2,3-dihydroxy-, (2 α , 3 β , 5 α , 25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 2137-20-4 HCAPLUS

CN Spirostan-3,12-dione, (5 α ,25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 883721-07-1 HCAPLUS

CN 22-Isoallospirostan-3 α ,12-diol (5CI) (CA INDEX NAME)

Absolute stereochemistry.



L97 ANSWER 16 OF 16 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1941:18004 HCAPLUS Full-text

DN 35:18004

OREF 35:2899c-i,2900a-b

TI Sterols. CXV. Sapogenins. 44. Relation diosgenin and cholesterol

AU Marker, Russell E.; Turner, D. L.

SO Journal of the American Chemical Society (1941), 63, 767-71

CODEN: JACSAT; ISSN: 0002-7863

DT Journal
LA Unavailable


AB The assumption that the C skeleton of the side chain in the steroidal sapogenins is identical with that of cholesterol (I) has been based on the isolation of α -methylglutaric acid from the oxidation products of digitogenic acid and the occurrence of Me isohexyl ketone (?) in the reaction products of Se with sarsasapogenin; this assumption has now been proved by converting diosgenin (II) into I and addnl. evidence is afforded for the 5,6-position of the double bond in II. Reduction of 5 g. II with 150 g. amalgamated Zn in 500 cc. EtOH and 150 cc. concentrated HCl for 3 hrs. gives 3 g. of tetrahydrosapogenin (III), m. 178-9°; triacetate (IV), m. 119.5°; IV is saponified by EtOH-KOH to III; tribenzoate of III, m. 166-7°. Catalytic reduction with PtO₂ of III, using 3 atmospheric of H for 2 hrs., gives tetrahydrodigitogenin (3,16,27-trihydroxycholestane) (V), m. 195-7°; catalytic reduction of IV or acetylation of V gives the tri-Ac derivative of V, m. 67-8°, tribenzoate of V, m. 162°. Refluxing 4 g. of IV in 25 cc. C₆H₆ with 2 g. H₂SeO₃ in 75 cc. 97% AcOH for 1 hr., adding 5 g. AcOK and refluxing for 10 min. give after hydrolysis with EtOH-KOH 0.5 g. of a tetrahydroxycholestene, C₂₇H₄₆O₄, m. 196°, which is converted by refluxing 1 g. with 5 cc. concentrated HCl in 100 cc. EtOH into Δ^4 -3-keto-16,27-dihydroxycholestene, C₂₇H₄₄O₃, m. 163-4°. Refluxing 4 g. III and 12 cc. PBr₃ in 300 cc. C₆H₆ for 2 hrs., purification of the product by washing the ether solution with H₂O and Na₂CO₃ and refluxing the residue (4.7 g.) in 150 cc. AcOH with 600 mg. AcOK with final reduction with Na in PrOH give Δ^5 -cholestene, m. 89-91°, and I, separated by sublimation (80-100° and 120-40°). Oxidation of 25 g. diosgenin acetate (VI) with CrO₃ in AcOH at 50-3° gives 4.4 g. unchanged VI, 50 mg. of an acid, C₂₇H₄₀O₅, decompose 226°, and 7-ketodiosgenin acetate (VII), m. 197° [semicarbazone (VIII), decompose 282°]. The Wolff-Kishner reaction with VIII gives a small quantity of 3,5-dehydrodesoxytigogenin. VII with 15% EtOHKOH (15 min. on the steam bath) gives 3,5-dihydro-7-ketotigogenin, m. 197-8°. Addition of 170 cc. HCl during 2.5 hrs. to 3 g. 4-dehydrotigogenone (IX) and 100 g. Zn-Hg in 500 cc. EtOH at the b. p. gives 500 mg. of 4-dehydrodesoxytigogenin, m. 145.5-6°; it also is formed with unamalgamated Zn. IX (5 g.) on reduction with (iso-PrO)₃Al in iso-PrOH gives 2.5 g. of 3,5-dehydrodesoxytigogenin, m. 168-9°; catalytic reduction yields desoxytigogenin, m. 173°. II (3 g.) and 17 g. p-C₆H₄O₂ in 200 cc. PhMe, from which 50 cc. of the PhMe is removed in vacuo, treated with 5 g. (iso-PrO)₃Al and refluxed 1 hr. give 0.9 g. of 4,6-dehydrotigogenone, m. 205-7°, which is purified by filtration through Al₂O₃ and treatment with succinic anhydride and C₆H₅N to remove carbinols. II gives chlorodesoxydiosgenin, m. 211-13°; catalytic reduction in AcOH yields 3-chlorodesoxytigogenin (X), m. 204-7°; reaction of 5 g. tigogenin in 100 cc. CHCl₃ containing 5 g. CaCO₃ with 5 g. PCl₅ gives 2.8 g. of an isomer(?) (XI) of X, m. 210-12° (mixed m. p. with X, 189-204°). Refluxing 1.3 g. XI with 30 cc. quinoline for 1 hr. gives 350 mg. of 2-dehydrodesoxytigogenin, m. 163-6°. Refluxing 5 g. 4-dehydrotigogenone and 15 g. (iso-PrO)₃Al in 500 iso-PrOH for 6 hrs., distilling slowly for 24 hrs. and rapidly to 0.5 its volume, cooling, adding 300 cc. cold 8% MeOH-KOH and after 1 hr. pouring into H₂O, gives 1.1 g. of 4-dehydroepitigogenin, m. 208-10°; this is not precipitated by digitonin; refluxing with Ac₂O for 30 min. gives Δ^3 ,5-desoxytigogenin; the material from the digitonin precipitate gives a compound, m. 167-9°, which is dehydrated by heating in vacuo at 100° and then m. 125-37°.

IT 512-04-9, Diosgenin
(and derivs.)

RN 512-04-9 HCAPLUS


CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.




IT 512-04-9P, Diosgenin 6870-79-7P, Tigogenone, 4-dehydro-
 16653-68-2P, Tigogenone, 4,6-dehydro-
 RL: PREP (Preparation)
 (preparation of)
 RN 512-04-9 HCAPLUS
 CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.




RN 6870-79-7 HCAPLUS
 CN Spirost-4-en-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.



RN 16653-68-2 HCAPLUS
 CN Spirosta-4,6-dien-3-one, (25R)- (CA INDEX NAME)

Absolute stereochemistry.



=> => d all hitstr retable

L105 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2007 ACS on STN

AN 1942:37236 HCAPLUS Full-text

DN 36:37236

OREF 36:5828f-h

ED Entered STN: 16 Dec 2001

TI Sterols. CXLVII. Sapogenins. 61. The bio-reduction of steroids

AU Marker, Russell E.; Wagner, R. B.; Ulshafer, Paul R.

SO Journal of the American Chemical Society (1942), 64, 1653-5

CODEN: JACSAT; ISSN: 0002-7863

DT Journal

LA Unavailable

CC 10 (Organic Chemistry)

AB cf. C. A. 36, 4516.3. From the feces of a 10-kg. dog, fed a mixture of 150 g. of meat, 50 g. of pig brain and 3 g. of diosgenin (I) for 3 consecutive days, there were isolated 5.2 g. of I, 0.2 g. of epismilagenin (II) and 0.1 g. of smilagenin (III) (as acetate). Similarly, tigogenone gives tigogenin and epitigogenin and sarsa apogenone yield sarsasapogenin and episarsasapogenin. This and earlier results (C. A. 36, 3182.9) support the hypothesis of Schoenheimer (C. A. 29, 353.4) that there is a reversible biol. reaction of the type cholestenone cholesterol. δ^4 -Dehydrotigogenone may be reduced by 1 enzyme system to II and III and by another system to I. The fact that HO compds. of both α - and β -configuration are formed is contrary to earlier statements (C. A. 32, 7471.9) that reduction in vivo of 3-ketosteroids appears to give only a compds.

IT Sapogenins

Sapogenins

Sterols

IT Steroids

(bio-reduction of)

IT Animal organism

(steroid reduction in)

IT 470-07-5, Tigogenone 512-04-9, Diosgenin

639-96-3, Sarsasapogenone

(fate in animal organism)

IT 126-18-1P, Smilagenin 16653-88-6P, Epismilagenin

RL: PREP (Preparation)

(formation in animal organism from diosgenin)

IT 126-19-2P, Sarsasapogenin 470-03-1P, Episarsasapogenin

RL: PREP (Preparation)

(formation in animal organism from sarsasapogenone)


IT 77-60-1P, Tigogenin 6788-40-5P, Epitigogenin

RL: PREP (Preparation)

(formation in animal organism from tigogenone)


IT 470-07-5, Tigogenone 512-04-9, Diosgenin
 639-96-3, Sarsasapogenone
 (fate in animal organism)
 RN 470-07-5 HCAPLUS
 CN Spirostan-3-one, (5 α ,25R)- (CA INDEX NAME)

Absolute stereochemistry.




RN 512-04-9 HCAPLUS
 CN Spirost-5-en-3-ol, (3 β ,25R)- (CA INDEX NAME)

Absolute stereochemistry.



RN 639-96-3 HCAPLUS
 CN Spirostan-3-one, (5 β ,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 126-18-1P, Smilagenin 16653-88-6P, Epismilagenin
 RL: PREP (Preparation)

(formation in animal organism from diosgenin)

RN 126-18-1 HCAPLUS

CN Spirostan-3-ol, (3 β ,5 β ,25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 16653-88-6 HCAPLUS

CN Spirostan-3-ol, (3 α ,5 β ,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 126-19-2P, Sarsasapogenin 470-03-1P, Episarsasapogenin


RL: PREP (Preparation)

(formation in animal organism from sarsasapogenone)

RN 126-19-2 HCAPLUS

CN Spirostan-3-ol, (3 β ,5 β ,25S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 470-03-1 HCAPLUS

CN Spirostan-3-ol, (3 α , 5 β , 25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 77-60-1P, Tigogenin 6788-40-5P, Epitigogenin


RL: PREP (Preparation)

(formation in animal organism from tigogenone)

RN 77-60-1 HCAPLUS

CN Spirostan-3-ol, (3 β , 5 α , 25R)- (CA INDEX NAME)


Absolute stereochemistry.



RN 6788-40-5 HCAPLUS

CN Spirostan-3-ol, (3 α , 5 α , 25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> d his

(FILE 'HOME' ENTERED AT 06:50:07 ON 13 DEC 2007)
SET COST OFF

FILE 'REGISTRY' ENTERED AT 06:50:17 ON 13 DEC 2007
ACT NOBLE531/A

```

-----
L1          STR
L2          2639 SEA FILE=REGISTRY CSS FUL L1
-----
L3          STR L1
L4          10 S L3 CSS SAM SUB=L2
L5          134 S L3 CSS FUL SUB=L2
           SAV L5 NOBLE531D/A
L6          17 S L5 AND 5 BETA
L7          117 S L5 NOT L6
L8          131 S L5 NOT (T/ELS OR 14C#)
L9          STR L3
L10         13 S L9 CSS SAM SUB=L2
L11         351 S L9 CSS FUL SUB=L2
           SAV L11 NOBLE351E/A
L12         23 S L11 AND NC>=2
L13         328 S L11 NOT L12
L14         295 S L13 NOT ((D OR T)/ELS OR 11C# OR 13C# OR 14C# OR C11# OR C13#
L15         12 S L14 AND IDS/CI
L16         283 S L14 NOT L15
L17         3 S L16 AND NR>=7
L18         280 S L16 NOT L17
L19         STR L9
L20         14 S L19 CSS SAM SUB=L2
L21         STR L19
L22         16 S L21 CSS SAM SUB=L2
L23         324 S L21 CSS FUL SUB=L2
           SAV L23 NOBEL531F/A
L24         12 S L23 AND NC>=2
L25         64 S L23 AND NR>=7
L26         52 S L25 NOT L24
L27         299 S L23 NOT ((D OR T)/ELS OR 11C# OR 13C# OR 14C# OR C11# OR C13#
L28         235 S L27 NOT L24-L26
L29         STR L9
L30         2505 S L29 CSS FUL SUB=L2
L31         STR L9
L32         352 S L31 CSS FUL SUB=L30
L33         1 S L32 NOT L11
L34         STR L21
L35         339 S L34 CSS FUL SUB=L2
L36         15 S L35 NOT L32,L23
L37         14 S L36 NOT 14C
L38         4 S 126-18-1 OR 470-03-1 OR 16653-88-6 OR 126-19-2
L39         1 S 512-04-9
L40         1 S 6870-79-7

```

FILE 'HCAPLUS' ENTERED AT 07:24:48 ON 13 DEC 2007

```

L41         2288 S L39 OR DIOSGENIN
L42         91 S L40 OR DIOSGENONE
L43         57 S L41 AND L42

```

```

L44      18 S L43 AND (REDUC? OR REDOX)
          E REDOX/CT
          E E34+ALL
L45      27061 S E9,E10,E11,E17
          E E9
          E E11+ALL
L46      139018 S E2-E4,E34,E35,E42,E43
          E E34+ALL
L47      113307 S E3-E5
L48      2 S L43 AND L45-L47
L49      16 S L44 NOT L48
L50      16 S L44 AND PY<=2002 NOT P/DT
L51      2 S L44 AND (PD<=20021028 OR PRD<=20021028 OR AD<=20021028) AND P
L52      2 S L48 AND L50,L51
L53      16 S L50,L51 NOT L52
          SEL AN 2 12
L54      2 S E1-E4 AND L53
L55      4 S L48,L54
L56      4 S L55 AND L41-L55
L57      2 S L56 AND (?SARSASAPOGENIN? OR ?EPISARSASAPOGENIN? OR ?SMILAGEN
L58      2 S L56 AND L38
L59      2 S L57,L58
L60      4 S L56,L59
          SEL RN

```

FILE 'REGISTRY' ENTERED AT 07:37:19 ON 13 DEC 2007

```

L61      30 S E5-E34
L62      8 S L61 AND (B OR AL)/ELS
L63      4 S L61 AND L8
L64      5 S L61 AND L18,L38
L65      6 S L61 AND L38,L39,L40
L66      2 S L61 AND L26,L28,L37
L67      11 S L61 NOT L62-L66

```

FILE 'HCAPLUS' ENTERED AT 07:38:18 ON 13 DEC 2007

```

L68      4 S L62-L66 AND L60
L69      1 S L68 AND LIALH4
L70      1 S L68 AND AL2O3
L71      4 S L68-L70

```

FILE 'REGISTRY' ENTERED AT 07:39:32 ON 13 DEC 2007

FILE 'HCAPLUS' ENTERED AT 07:40:03 ON 13 DEC 2007

```

L72      297 S L8
L73      193 S L72 AND (L18 OR L38 OR L28 OR L26 OR L37)
L74      164 S L73 AND PY<=2002 NOT P/DT
L75      15 S L73 AND (PD<=20021028 OR PRD<=20021028 OR AD<=20021028) AND P
L76      179 S L74,L75
L77      3 S L76 AND L45-L47
L78      0 S L76 AND REDOX
L79      48 S L76 AND REDUC?
L80      48 S L77,L79
L81      1 S L80 AND L62
L82      22 S L80 AND (LIALH4 OR AL2O3 OR ?BORON? OR ?BORAN? OR ?BORIC? OR
L83      22 S L81,L82
L84      46 S L80 NOT L71

```

FILE 'REGISTRY' ENTERED AT 07:42:33 ON 13 DEC 2007

FILE 'HCAPLUS' ENTERED AT 07:42:33 ON 13 DEC 2007

L85 TRA L84 1- RN : 1015 TERMS

FILE 'REGISTRY' ENTERED AT 07:42:35 ON 13 DEC 2007

L86 1015 SEA L85
L87 1 S L86 AND (B OR AL)/ELS
L88 1 S L86 AND (BORON? OR BORAT? OR BORIC? OR ?ALUMIN?/CNS)
L89 1 S L87,L88

FILE 'HCAPLUS' ENTERED AT 07:43:22 ON 13 DEC 2007

L90 1 S L89 AND L80
L91 22 S L83,L90
L92 18 S L91 AND (L18 OR L38 OR L28 OR L26 OR L37) (L)PREP+NT/RL
L93 3 S L91 AND L8 (L) RACT+NT/RL
L94 2 S L92 AND L93
L95 1 S L94 NOT L71
L96 17 S L92,L93 NOT L94,L71
L97 16 S L96 AND L18,L38
L98 20 S L83 NOT L71,L95
L99 1 S L98 AND L8(L)RACT+NT/RL
L100 16 S L98 AND (L18 OR L38 OR L28 OR L26 OR L37) (L)PREP+NT/RL
L101 0 S L99 AND L100
L102 25 S L80 NOT L71,L95,L100,L98
 SEL AN 21
L103 1 S L102 AND E35-E36
L104 2 S L99,L103
L105 1 S L104 NOT L71,L95,L97

=>